

6th World Congress and Expo on

Breast Pathology and Cancer Diagnosis

&

20th International Conference on

Medicinal Chemistry and Rational Drugs

July 25-26, 2018 | Vancouver, Canada

Study of the iron chelating effect of green tea in smear positive TB patients using sputum smear, serum malondialdehyde and blood iron indices

Shahryar Eghtesadi, Honarvar M R, Gill P, Jazayeri S, Vakili M A, Shamsardakani M, Abbasi Azad University, Iran

Green tea with possessing iron chelating properties can be useful in TB treatment and management. We studied the effect of green tea consumption on iron status and improving process of pulmonary tuberculosis treatment (accelerating the negative sputum smear, reducing the level of oxidative stress). Following the approval by Ethics Committee for Human Studies Golestan and Tehran Universities of Medical Sciences and also obtaining the written consent of patients, this double-blind randomized clinical trial study, was conducted on patients with TB, who were assigned randomly to the intervention group (41 patients) receiving 500 mg catechin of green tea extract and the control group (39 subjects) receiving placebo for two months, since the beginning of concomitant anti-TB treatment. Sputum evaluation was carried out on three slides using the Ziehl Nelson method. At first, the demographic and dietary intake data were obtained. After obtaining 10 ml of venous blood, Hemoglobin (Hb), Transferrin, Ferritin, Total iron binding capacity (TIBC), Iron and Serum malondialdehyde (MDA) were measured at the beginning and end of the study. Sputum samples were collected from the third week (every 10 days) and the reduction of microbial load was also tested until sputum smear became negative. Data were processed using independent and paired t-test, McNemar, Wilcoxon, Kaplan-Meier, Log-rank test and Cox regression model. P-value was taken significant as <0.05. Average daily energy intake of patients was 1518±431 kcal, distribution of which was as follow: carbohydrates (58%), protein (17%) and fat (22%). Vitamin D and Zinc intake of patients were less and iron intake was higher than the DRI. Weight changes in both groups of placebo and green tea had tendency of increase with a significant difference at two and six month follow ups (p>0.0001). However, there were no significant changes due to intervention compared to placebo. Sputum conversion time (days) was 52.5± 24.5 (median= 53 days) and 40.6 ± 22.5 (median= 29 days) in placebo and catechin groups, respectively. Conversion time of patients in the green tea group based on criterion of; the short duration of being negative sputum smears was significantly higher than the placebo group (p=0.032). To measure the mean of iron status after intervention, ANCOVA test showed mean difference level (Pvalue) in both groups for Hb, iron, TIBC, transferrin and ferritin as of: 0.004, 0.56, 0.65, 0.38 and 0.16, respectively which means that increase of hemoglobin in the green tea group was significant compared with the placebo group. There was just a 9.2 nmol/ml difference between the two groups for MDA at the beginning of study, which was not statistically significant (p=0.078) whereas, it was increased to 24.8 nmol/ml after the intervention, indicating a significant difference (p<0.001). The decline value was estimated -45.45 ± 14.69 nmol/ml for catechin group and -19.91 ± 18.38 nmol/ml for placebo group. In conclusion green tea can systematically reduce the inflammatory elements and oxidants (decrease of MDA as fatty acids oxidation indicator), and consequently, can improve the hematopoiesis and hemoglobin level. Therefore, localized inflammation and damage in the lung is reduced, and adjunct to antimicrobial therapy, accelerate sputum smear conversion, disease amelioration and treatment improvement. Finally, given the higher iron intake despite of lower micronutrients and macronutrients in diet of our patients, and considering the iron effect on mycobacterium survival and the incidence and exacerbation of inflammatory complications in patients, it seems that policy of mandatory iron fortification with iron, especially in provinces such as Golestan, must be viewed cautiously and its further implementation being revised meticulously.

Biography

6 KDKU\DU (JKWHV DGL UHFHLYHG %DFKHORU GHJUHH LQ 1XWULWLRQ 6FLHQFH DQG)RRG &KHPLVWU\ GHJUHH LQ 1XWULWLRQ IURP 7HKUDQ 8QLYHUVLW\ RI OHGLFDO 6FLHQFHV 7HKUDQ DQG 3K' IURP 8 VHUHG DV 9LVLWLQJ 6FLHQWLWV LQ 86'\$ +XPDQ 1XWULWLRQ 5HVHDUFK &HQWHU RQ \$JLQJ +15&\$ % 8QLYHUVLWLV RI OHGLFDO 6FLHQFHV DQG FXUUHQWO\ VHUHG DV 3URIHVVRU RI \$JG 8QLYHUVLW\ 6F DQG %LRFKHPLVWU\ %LRFKHPLVWU\ &OLQLFDO 1XWULWLRQ 3XEOLF +HDWK 1XWULWLRQ DQG 1XWUL RI 6FKRRO RI 3XEOLF +HDWK 1XWULWLRQ DQG 6FKRRO RI 3XEOLF +HDWK RI 7DEUL] DQG ,UDQ 8QLYH SURIHVVRU DQG 6FLHQWLWV)RU ORQJ DQG H[WHQGHG SHULRG RI WLPH H[SHULHQFHG WHDFKLQJ YDU LQWHUQDWLRQDO %XUHDX SURJUDPV DQG GLUHFHWG PDQ\ SURMHFWV DQG GLVHUWDWLRQ RI 06 DQG DQG DOVR HGLWHG VHYHUDO ERRNV DQG QDROD &HWDQHG DUJRU SHULQV ILSD & DSHFFLW L JDXLWLG IRQJ LQ 1XWUL

VHJWHVDGL#JPDLO FR