**h**tDg St2017, 7:4(S**p** DOI: 10.4172/2161-069X-C1-053

## 12TH EURO-GLOBAL GASTROENTEROLOGY CONFERENCE

September 11-12, 2017 | Paris, France

Rose hip in treatment of non-alcoholic fatty liver disease (NAFLD)

Hadis Sabour Tehran University of Medical Sciences, Iran

Non-alcoholic fatty liver disease (NAFLD) is one of the most common liver diseases in Iran. e disorder is being recognized increasingly in the uniformly obese and no obese people. However, the lack of speci c reversal agents has made the management of (NAFLD). Rose hip has been used traditionally to treat a range of conditions including diabetes based on anti-insulin e ect of rosehip. is study on 25 patients illustrates the role of rosehip in improving patient Non-alcoholic fatty liver disease while highlighting the importance of information and education about the availability and appropriate use of this herb in treatment of fatty liver.

## Biography

Hadi Sabiri cerhig a: AibtPf. iBarad SpillinReeach Cebr(BASIR), Neirceoe Ibe, Tehari Jiatif Medical Scene: (2016 t date). He ha: iled a: Reeacher Det Crist Barad SpillinReeach Cebr(BASIR), Neirceoe Ibe, Tehan Ubatif Medical Scene: (2006-t2015) ad Reeacher Edicia ad Meabli Reeach Cebr Tehari Jiatif Medical Scene: (2005-2007). He ha: ciped hi PhD (2011) in Ital Scene: fr Tehari Jiatif Medical Scene:, Teharad MD i 2000 fr Tabiz Jiatif Medical Scene: Scene: Tehari Jiatif Medical Scene: Tehari Jiatif Medical

habi@ia.t.ac.ir

**Notes:**