

4<sup>th</sup> International Conference on

# Palliative Care, Medicine and Hospice Nursing

August 27-28, 2018 | Boston, USA

## Safety of wearing contact lenses for patients using selective serotonin reuptake inhibitors (SSRIs)

Yi-Pei LIU, Yi-Jiun LIU, and Yi Shuan LIU  
Taipei City Hospital, Taiwan

**Objective:** To evaluate the safety of wearing contact lenses for patients using selective serotonin reuptake inhibitors (SSRIs). The study included 100 patients who were wearing contact lenses and were prescribed SSRIs. The patients were divided into two groups: the control group (50 patients) and the SSRI group (50 patients). The control group wore contact lenses for 2 weeks, and the SSRI group wore contact lenses for 2 weeks. The safety of wearing contact lenses was evaluated by the number of patients who experienced contact lens-related complications (CLRCs) during the 2-week period. The results showed that the number of CLRCs was significantly higher in the SSRI group than in the control group.

**Methods:** A retrospective study was conducted to evaluate the safety of wearing contact lenses for patients using selective serotonin reuptake inhibitors (SSRIs). The study included 100 patients who were wearing contact lenses and were prescribed SSRIs. The patients were divided into two groups: the control group (50 patients) and the SSRI group (50 patients). The control group wore contact lenses for 2 weeks, and the SSRI group wore contact lenses for 2 weeks. The safety of wearing contact lenses was evaluated by the number of patients who experienced contact lens-related complications (CLRCs) during the 2-week period.

**Notes:**