

14th World Congress on

TOXICOLOGY AND PHARMACOLOGY


March 12-14, 2018 Singapore

Development of a screening kit for detecting synthetic cathinones

Yuki Mukai, Fujio Ishizawa, Hiroko Masuda, Hisanori Muramatsu and Katsuya Honda
University of Tsukuba, Japan

In recent years, abuse of illicit drug has been a very important problem.

Significant amount of unknown illicit substances are sized by enforcement and boarder protection agencies. Synthetic cannabinoid cathinones have a psychoactive effect on our bodies. The identification of these drugs is important not only for the proof of the crime, but also human



health. Generally, to identify the illicit drugs, we use the techniques such as gas chromatography-mass spectrometry (GC-MS) and high performance liquid chromatography-mass spectrometry (HPLC-MS). These instruments, however, are not always convenient owing to their high cost of running, the need for trained personnel, lengthy analysis times, etc. Screening kits for detecting drugs, therefore, are required at the scene of a crime and the development of them is desired. In this study, we have designed the screening kit for cathinones to provide easy indication of the presence or absence of cathinones in a test sample. This kit consists of a glass tube enclosed reagents which react with them (named "cathinone detector tube"). This kit utilizes the reaction of cathinons with neocuproine and copper(II) to give a colored copper(I)-neocuproine complex. The presumptive color test method for the detection of synthetic cathinones by Morgan Philip, et al. is applied to the development of this screening kit. They describe that neocuproine color test displays good selectivity to cathinone analogs. To improve operativity and preservation of our kit, three aqueous solutions were coated on silica gel particle and then the powders have been enclosed in a single glass tube. The reagent in the tube colored orange from light blue in the presence of cathinones. In consequence, this kit had very high sensitivity for detecting cathinones. The limit of detection of -PVP, for example, was 5 µg (absolute amount, 100 µL of 50 µg/mL solution).

References

1. Morgan Philip, Ronald Shimmon, Mark Tahtouh, Shanlin Fu (2016) Development and validation of a presumptive color test method for the detection synthetic cathinones in seized illicit materials. *Forensic Chemistry*: 39-50.

Biography

<XNL 0XNDL LV FXUUHQWO\ D 0DVVHUV VWXGHQW DW 8QLYHUVLW\ RI 7VXNXED -DSDQ 6KH EHORQJV WF
SUREOHP LQ WKH ZRUOG HVSHFLDOO\ LQ V\QWKHWLF FDQQDELQRLGV DQG FDWKLQRQHV 6KH KDV UHF

V #X WVXNXED DF MS

Notes: