

6th World Conference on Climate Change

September 02-03, 2019 | Berlin, Germany

Long-term effects of residential rainfall levels on post heavy-rainfall waterborne disease hospitalizations in young children in New Zealand

Hakkan Lai, Caroline Walker, Alistair Woodward, Peter Tricker and Susan Morton
University of Auckland, New Zealand

Climate change models have recently projected increased heavy rainfall events in both wet and dry regions. It is well established that heavy rainfall events are associated with increased risks of waterborne diseases (WD). However, there is a paucity of epidemiological evidence for how wet and dry regions influence the risk of WD due to heavy rainfall. In New Zealand, the annual total rainfall can differ by up to 3500mm in different locations. We aimed to determine if residing in wet or dry regions was associated with the risk of experiencing WD immediately following heavy rainfall. Using our cohort of 6853 children born between 2009 and mid-2010 in New Zealand, we determined the possible WD hospitalization if the primary and/or secondary causes were intestinal infections (ICD10:A00-09), non-intestinal E. coli infections (B96.2), leptospirosis (A27), mycobacterial infections (A31), adenovirus (B97.0), enterovirus (B97.1), and/or unspecified-site viral infections (B34). We only considered admissions within a lag window of 1-4 days after the heavy rainfall dates to ensure short-term temporality. To determine wet or dry regions, we assessed time-weighted long-term exposure using median annual rainfall levels at individual home locations at various different data collection time points. We used logistic regression model adjusted for child's sex, ethnicity, rurality and individual deprivation. Based on the third quintile of residential long-term rainfall level, the adjusted odds ratios [95%CI] of post heavy-rainfall WD hospitalizations in the dry (first, second) and wet regions (fourth and fifth quintiles) were 1.84 [1.08-3.14], 1.23 [0.70-2.17], 1.35 [0.77-2.37] and 2.24 [1.25-4.01] respectively. A U-shaped exposure-response relationship was found (quadratic trend P-value = 0.002). Living in the wettest and driest rainfall locations were both associated with childhood WD hospitalizations shortly after heavy rainfall dates. We suggest a need to review preventive policy to address the rainfall-associated WD risks among residents in vulnerable locations.

Recent Publications

1. Lai, H. K., Camargo, C. A., Woodward, A., Hobbs, M., Pillai, A., Morton, S. M., Grant, C. C. (2019). Long-term exposure to neighborhood smoke from household heating and risk of respiratory and dermatological prescription medications - Growing Up in New Zealand child cohort study. *Allergy* 74:391-395.
2. Morton, S. M., Lai, H. K. (2018). Growing Up in New Zealand: Health research project – Drinking water quality and child health. In *Water New Zealand Conference & Expo 2018*. Hamilton, New Zealand.
3. Lai, H. K., Berry, S. D., Verbiest, M. E., Tricker, P. J., Atatoa Carr, P. E., Morton, S. M., & Grant, C. C. (2018). Emergency department visits of young children and long-term exposure to neighbourhood smoke from household heating - the Growing Up in New Zealand child cohort study. *Environmental Pollution* 231(Pt 1), 533-540.

Biography

%HIRUH VWDUWLQJ SRVW GRF UHVHDUFK LQ HQYLURQPHQWDO KHDOWK LQ +RQJ .RQJ 'U +DNNDQ /D .Q KH PRYHG WR 1HZ =HDODQG DQG LQ KH MRLQHG WKH 8QLYHUVLW\ RI \$XFNODQG 6FKRRO %LRVWDWLVLWLFV IRU D VWXG\ UHODWHG WR ELRGLYHUVLW\ JUHHQ VSDFH DQG DFWLYH WUDQVSRU +DNNDQ¶V UHVHDUFK IRFXVHV RQ VSDWLDO DQG WHPSRUDO GLYWULEXWLRQ RI HQYLURQPHQWDO H respiratory, gastrointestinal, dermatological and oncological outcomes