


Commonwealth of Saint Kitts & Nevis Orthodontist & Dentofacial  
Orthopedist



oapp• Closed orthodontic eruption.

However, not all impacted teeth are available to be treated by  
the enforced orthodontic eruption and not all the impacted teeth

1- Extraction of the impacted tooth with poor prognosis or in the





DDS, MSC









Clinical choice of the most appropriate implant therapy modality should be based on assessment of the residual alveolar bone height, width, and sinus morphology with a cone beam

preferences. Following a good surgical protocol and excellent oral hygiene maintenance program are fundamental elements in achieving a successful and predictable outcome. The available evidence on short dental implants in early research was not sig-

treatment is improving now than before for this reason.

The use of short implants allows treatment of patients who are unable to undergo complex surgical techniques for medical, ana-

geries short implants reduce morbidity, cost and treatment time.

Recently short implants offer a less invasive treatment alternative in resorbed ridge cases.

tistry. Head of the Dental Department and Dental Educator, Director of Academic and Education Affairs at Comprehensive

ical Services, Jeddah, Kingdom of Saudi Arabia, where he has been since 2016. He received a Saudi Fellowship In Dental Implant from the Saudi Commission for Health Specialties, 2014-2016.

He received a Saudi Board In Periodontics from the Saudi Commission for Health Specialties, 2012 -2014.

He received his Master of Science in Dentistry (MSD) and a Clin-

and Pharmacy 2009-2012, Riyadh, KSA. With Honours. Also he received his Advanced Education in General Dentistry (AEGD)

gery (BDS) from Faculty of Dental Medicine and Surgery, King