

Tricuspid Stenosis due to Cardiac Sarcoma

Mirza Baig
Gwynedd Hospital, UK

Abstract

Primary cardiac sarcomas are a rare and aggressive tumour type with a high rate of recurrence, irrespective of treatment, and a poor prognosis. The age of presentation for cardiac sarcomas ranges from 1 to 76 years, with a mean age of around 40 years. Angiosarcoma

cardiac sarcomas, of which angiosarcomas are the most common. Angiosarcomas are predominantly found on the right side of the heart. Pericardial

Here we present a 32- year old male with Asian ethnicity, symptoms including-abdominal pain, nausea and vomiting. On initial evaluation, everything pointed toward a gastroenterology (GI) aetiology, but increasing signs of right heart failure and an

cardiac surgical team and to local oncology once the diagnosis of primary cardiac sarcoma had been made. He has gone on to awaiting oncology and cardiology review. Although the diagnosis of primary cardiac sarcoma is rare, it should be considered in

has thus been recently cardioverted. Presently, he is doing well and

DQG XQFODVVL¿HG VDUFRPDV DFFRXQW IRU DSSUR[LDPWHO\ RI DOO WKH KHDUW ZKHUHDV RVWHRVDUFRPDV DQG XQFODVVL¿HG VDUFRPDV DUH DQJLRVDUFRPD LV H[WUHPHO\ UDUH

UHFHQWO\ PRYHG WR WKH 8. ZLWK D KLVWRU\ RI QRQ VSHFL¿F LQWR KLK ULJKW YHQWULFOH [FP LQ VL]H ZLWK DQ DVVRFLDWHG SHULFDUGLDO HuXVLRQ +H ZDV VXEVTXHQQWO\ UHIHUHG WR ERWK WKH GHYHORS DWULDO AXWWHU DQG VHYHUH OHIW YHQWULFXODU G\IXQFWLRQ DQ \RXQJ SDWLHQWV SUHVHQWLQJ ZLWK QRQ VSHFL¿F \PSWRPV DQG VLJQV