



A Retrospective Chart Analysis of the Experiences of Breast Cancer Patients Using Telehealth

Raouf Hano*

Department of Human Physiology, College of Colchester, United Kingdom

Abstract

The most common malignant tumour in women is breast cancer. Several approaches are used in its therapy. In remote areas, telehealth can provide continuity of care. The goal of this study was to compile qualitative data on

-
- stearothermophilus unfolds through a monomeric intermediate. A quantitative analysis under equilibrium conditions. *The J Biol Chem* 273:18052–18059.
5. Ould-Abeih MB, Petit-Topin I, Zidane N, Baron B, Bedouelle H (2012) Multiple folding states and disorder of ribosomal protein SA, a membrane receptor for laminin, anticarcinogens, and pathogens. *Biochemistry*. 51:4807–4821.
 6. Cahn RS, Ingold CK, Prelog V (1966) Specification of Molecular Chirality. *Angew Chem Int Ed* 5:385–415.
 7. Vickery HB, Schmidt CL (1931) The history of the discovery of the amino acids. *Chem Rev* 9:169–318.
 8. Ntountoumi C, Vlastaridis P, Mossialos D, Stathopoulos C, Iliopoulos I, et al. (2019) Low complexity regions in the proteins of prokaryotes perform important functional roles and are highly conserved. *Nucleic Acids Res* 47:9998–10009.
 9. Marcotte EM, Pellegrini M, Yeates TO, Eisenberg D (1999) A census of protein repeats. *J Mol Biol* 293:151–160.
 10. Magee T, Seabra MC (2005) Fatty acylation and prenylation of proteins: what's hot in fat. *Curr Opin Cell Biol* 17:190–196.