

A Short Commentary about Efficacy of an Flowable Matrix in the Treatment of Diabetic Foot Ulcers

Campitiello F, Mancone M^{*}, Della Corte A, Guerniero R and Canonico S

Department of Medical, Surgical, Neurologic, Metabolic and Aging Sciences, University of Campania "Luigi Vanvitelli", Italy

***Corresponding author:** Mancone M, Department of Medical, Surgical, Neurologic, Metabolic and Aging Sciences, University of Campania "Luigi Vanvitelli", Italy, Tel: 3388212431; E-mail: manfredi.mancone@alice.it

Rec date: Jul 06, 2017; **Acc date:** Jul 22, 2017; **Pub date:** Jul 28, 2017

Copyright: © 2017 Campitiello F, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The authors aimed to evaluate the efficacy of an advanced wound matrix (Integra Flowable Wound Matrix, Integra

After severe trauma, in the adult mammal occurs a wound constriction and scar formation. An appropriate control of wound contraction can induce the regeneration of wounded skin [20]. More specifically it has been observed that the regeneration of the highest quality is induced under wound conditions the contraction was

17. Stiefel D, Schiestl C, Meuli M (2010) Integra artificial skin for burn scar revision in adolescents and children. *Burns* 36: 114–120
18. Weigert R, Choughri H, Casoli V (2011) Management of severe hand