

Adenocarcinoma Arising from Fistula in Ano

* Fazl Q. Parry, SKIMS, Colorectal surgery, 44 rawalpora, govt housing colony, sanat nagar, srinagar, Jammu and Kashmir, 190005, India, Tel: 9419008550; E-mail: fazlparry@rediff mail.com

Oct 2, 2015, Dec 1, 2015, Dec 7, 2015

© 2015 Nazki S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Adenocarcinoma

Previously reported adenocarcinomas are of low grade in about 44% percent cases [12]. One of the study concluded the final management as abdominoperineal resection with wide tumor excision on the gluteal region and reconstruction by bilateral gracilis musculocutaneous flaps [13]. It is important to look for unusual changes like persistent discharge, gradually increasing perianal mass, bloody discharge, unusual healing or induration in case of perianal fistula. Biopsy should be taken in all cases to be on the safer side.

1. Kyzer S, Bayer I, Turani H, Chaimov C et al. (1985) Verrucous squamous carcinoma as a complication of recurrent multiple perianal fistulae. *Coloproctology* 7: 104-106
2. Salat A (2006) Anal cancer EurSurg 38: 135-138
3. Heidenreich A, Collarini HA, Paladino AM, Fernandez JM, Calvo TO (1986) Cancer in anal fistulas: report of two cases. *Dis Colon Rectum* 9: 371-376
4. P