

Appearances of Stigma and Discrimination among Immunocompromised Female Injecting Drug Users (Fidu) – A Study in Champai, Mizoram in India

¹Department of Virology, ICMR National Institute of Cholera and Enteric Diseases, Kolkata, India
²Community Care Center, Champai, Mizoram, India

In India the HIV positivity in among IDUs stands at a staggering 7.71. Injecting drug use among female appear to mirror patterns among males, but with greater adverse consequences. They are still a group of population that lacks visibility, and are subjected to multiple layers of stigma because they belong to socially deviant and disenfranchised *! [~] • Á, ñc@h-æ&â } * Á * ^) ä ! É •] ^ & ä, & ñ } ^ ~ æ j ñ c ^ ñ æ } ä ! ^ c & j ~ • ä [] É Á V @ ^ h • c ~ ä ^ ñ æ ä { ^ ä ñ c h ~ } ä ! ^ • c æ } ä ä } * h c @ ^ Á, æ ^ • ñ } Á, @ ñ & @ h 0 Ö W • Á Ö @ æ {] æ ñ ä ä • c ñ ä & c Á [- Á T ä : [ñ æ { ñ ä } ä ä * ^) [~ • h { ä } [ñ ä c ^ h & [{ { ~ } ñ c ^ É Á c } ^ ñ ä } & ^ h • c ä * { æ ñ æ } ä ä ä ä • ñ ä { ä } æ c ä [] æ } ä h c @ ^ h {] æ & c • h c @ æ c h stigma and discrimination may have on their abilities to access health services. The qualitative study also used content analysis and the health stigma and discrimination framework analysis. The study found several forms of stigma among - ^ { æ j ^ h ä } b ^ & c ä } * ä ä ! ~ * ä • h c @ æ c h ä c ^ h • h c @ ^ ä ä æ & ^ • h æ } ä h ^ c ä ä : æ c ä [] h [- Á 0 æ j c @ h • h c ä } & ^ h • h æ } ä h • * * ^ c ^ ä h æ ä ! æ } * ^ h @ ^ æ j c ^ U c ä ^ æ ^ ä ! • \

Jan-2022, Pre QC No: dementia-21-35299 (PQ), 18-Jan-2022, QC No: dementia-21-35299, 24-Jan-2022, Manuscript No: dementia-21-35299 (R), 31-Jan-2022, DOI: 10.4172/dementia.1000117

study on diseases and identities of HIV.

Ghosh GK, Vanlalhumi (2022) Appearances of Stigma and Discrimination among Female Injecting Drug Users (FIDUs), they belong to socially deviant and disenfranchised groups with facing gender-specic inequality, stigma and exclusion in Champai, Mizoram in India. J Dement 6: 117.

© 2022 Ghosh GK, et al.

poor social support characterize this group. Sexual and reproductive health problems are frequent, particularly abortions. Even though evidences suggest an increase in the number of Female Injecting Drug Users (FIDUs), they belong to socially deviant and disenfranchised groups with facing gender-specic inequality, stigma and exclusion in Champai, Mizoram in India. The study attempted to focus on FIDUs of Mizoram state, Champai district, situated at Myanmar border, known for the illegal drug route. Mizoram estimated more than 28 thousand people injecting drugs for non-medical purposes –highest among all states. The HIV prevalence of 19.8% was considered stable. Heroin addiction among young people of both genders is 81.7% and injecting drug use affected 96.2% young people of Champai district with 61.2% sharing of injecting drug use reported for the district [7].

Method

A qualitative study was conducted in August and September 2019 among Female HIV positive intravenous drug users registered in the antiretroviral therapy centres (ARTC) for more than two years. Data for qualitative study were collected through focus groups with 14 HIV positive FIDUs, conducted in local Mizo twang dialect, were semi-structured consisting of a series of broad open-ended questions. Five individual interviews with ART service providers and stakeholders were undertaken to obtain their 'views on FIDUs' experiences and perspectives. The study used content analysis based on the health stigma and discrimination framework, as proposed by Strandberg, Parshlow, et al [8]. We embarked at finding out the antithesis fallout on a selected population i.e. stigmatized persons or groups, as well as their family and/or healthcare providers. Accordingly, we focused on conceptualizing the study on diseases and identities of HIV, taking cue from the foresaid study, to construct relevant health stigma and discrimination framework for the study, as in Figure 1 below.

Settings: The focus groups were conducted at participants' convenient location near to New Hope Society- care & support centre organization (CSCO). All the participants were linked to CSCO, but none of reportedly linked to any intravenous drug user (IDU) targeted intervention (TI) and not covered under needle-syringe exchange programme (NSEP), nor were they receiving opioid substitution

Objective

The study was conducted among hard-to-reach female injecting drug users with the view to understanding the impediments to their proper healthcare service uptakes, especially reproductive health care and harm reduction services.

Hypotheses

The study embarked at finding out the antithesis fallout on a selected population

i.e. stigmatized persons or groups, as well as their family, friend or healthcare providers. Accordingly, we focused on conceptualizing the

therapy (OST). However, they received CSCO services as counselling, periodic CD4 count test guide, and referrals, as stated.

Participant selection and recruitment: In the current study a person who has injected at least once in the last three months is categorized as an IDU in keeping with the definition followed in the National AIDS Control Programme (NACO). She should be above 18 years of age and give consent to participate.

Sampling: Given the hidden nature of FIDUs, we attempted snowball sampling to recruit participants for the study. Altogether 18 FIDUs were approached at the ARTC among whom 2 females declined. Only 14 among whom 10 were males and 4 were females participated.

8 said, 'I usually don't like people to know my drug injecting status because I feel shame and when going out invariably use body covered clothes to hide injection marks'.

Experienced stigma- Many participants stated to have experienced negative attitude towards injecting drug user from their communities. 'When I started doing drugs and injecting, others who do not use, refused to mix with me. I was able only to associate only with fellow drug users' – informed FGD participant 5. Two other FGD respondents also stated they were also stigmatized by some of their family members: 'From my own perspective, even the CSC workers are more concerned with us and love us even more than our relatives'- respondent 9 stated. And respondent 11 added, 'If you try to reason out or explain something in the family settings, they brush you aside as an addict. It pains. "Among the participants 42% females were earning either from petty business or in some job. Stigma and discrimination at workplace seemed experienced by some. Said FGD participant 12, a 28 years old single female, 'I was working in a beauty parlour. Once my HIV positive status became known to the owner, he instructed that I leave the job in the interest of the parlor and find engagement somewhere. I felt devoted'. Participants described how this stigma was invoked within a milieu of social dynamics and moralistic attitudes, conveying a notion of moral judgments and suspicion: 'People stated to me that a ruihhlo ngai (drug user) is a sinner'- FGD respondent 14 stated. Another respondent 7, informed, 'My school-day friends and neighbors become unduly alarmed by my presence near to their houses, as if I am thief. Very demeaning for Me.' Apart from experiencing stigma and discrimination in public spaces, FIDUs frequently experienced the same in healthcare related settings. Respondent 10 reported 'health care staff at the centre have a queer look at us, and whisper among fellow staff on my doing drugs and injecting habits'. Some participants also lamented over their non-existent access to needle syringe exchange programme and opioid substitution therapy under the IDU TIs.

Stigma of being FIDU was layered over HIV-related stigma, as respondent 11 described preceding HIV stigma, even from their fellow drug users. According to FGD respondent 3, 'Once I was detected HIV positive through community based testing; my fellow drug user asked me not to come in their group'. This statement suggested that being HIV positive was different source of shame among women who inject drugs.

Given these deeply ingrained cultural beliefs and taboos, coping with multi-layered stigma is particularly challenging for females. Despite this, PLHIV FIDUs found to hold positive internal beliefs about themselves as- 'I have changed over the years. It's like, I don't care who knows, everybody can know,' respondent 9, a 36 years old lady, a separated from marriage and in a government, on ART for the last seven years asserted. Still another respondent-13, a 26 years old single lady, with college level education, on ART regimen for 5 years and doing petty business, stated- 'I have understood that worrying won't help. I take my ART medicines regularly and keep going.'

Negative impact of stigma- Notwithstanding the source, stigmatization of women seemed to result in isolation and exclusion within the community. 'I feel isolated from my friends and family members.'

they said 'HIV/AIDS specialists treated us normally and quite helpful to PLHIV without being judgmental toward IDUs'.

Internalized stigma- participants' account indicated a self-directed shameful consciousness of being an injecting drug user female and PLHIV. Internalization of stigma was often associated with low self-esteem among participants as- 'it is difficult to tell others of my HIV positive status, being IDU' – FGD participant 2 stated. 'I feel worthless being IDU and also as HIV positive' – another FGD participant 13 added. Both the statements carry participant's negative reaction about self in varying degrees. Self-stigmatization resulted from study participants internalizing the negative perception of injecting drug use within the study setting, where drug use was generally construed as resulting from a failure of individual morality. FIDUs were commonly referred as 'hmeichhia ruihhlo ngai' (female drug user) and they are often thought of as 'K.S.' (slang word for female sex workers). Not surprisingly, FIDUs found such identities shameful and they had an interest in concealing their stigmatized identities. As FGD respondent

health service access. Participants characterized the actions of health system functionaries 'disdainful maltreatment' as FGD respondent 2 commented. Another FGD respondent 7 stated, 'Should they know that one is an addict, they send the patient backward on the queue or tell the person to go and come later'.

Apart from the negative perception of drug use, CSC functionary pointed out- 'the health care workers did not understand why and how they need to serve female drug users. Nevertheless, given women's experiences, being stigmatized at health facilities meant that 'she had never gone back' he added. But, studies indicated that many women inevitably found themselves compelled by circumstances to seek health services, and in those situations, they strived to conceal their identities, so as to avoid being stigmatized or discriminated against [11]. In Champai many participants opted to be accompanied by CSC outreach workers to the health centers, to avoid being feeling embarrassed. One FGD respondent-4 claimed, 'if we go to hospital with the out-reach workers (ORW), the health providers give us the required services without asking many question'

Discussion

In review of relevant documents and published research articles, the study endeavoured at analysing the various forms of stigma and discrimination faced by PLHIV FIDUs of Champai on the conceptualized stigma framework as in figure-1, to understand the stigma drivers and facilitators. Interactions with key informants indicated that lack of public awareness and knowledge on HIV issues concerning intravenous drug use, especially by females, existed, as the CSCO functionary said "*i a e d g e i c ide ed a i h HIV a i g heed he fac ha i jec i g d g, e e, d e ead HIV; i i he ha i gi fec ed i jec i ge i e ha d e . IDU i he egi e e f a e bjec ed a high e e f ig a a d d i c i i a i , a d b a ia ed fea fi fec i e i ed a, FIDU a e b a ed f he i fec i*". He also asserted - "*D g e a, a he, ed a a fa hi a d a fad f he gi he egi . e f a e i e f ch ig a he IDU ade he hide hei habi. e ed acce i g he e ice ided f hei afe a d e bei g*".

Notably, high literacy rate among females in Mizoram is recorded and all FIDUs participants at Champai held minimum of middle school education. Sharing of injecting paraphernalia by females with their male partner or spouse and unprotected sex were reportedly the dual modes of HIV transmission among FIDUs. In Champai, males continue to be main earning members for the family; and as such females have minimum say in economic matters, in decision making and obviously weak negotiating capacities in conjugal life. Again, among sensitized females, indulging in risky behavior under influence of drugs with male IDUs and partners found common, since IDU is considered by them 'enjoyable group activity'. Said the ARTC counsellor, "Females are discriminated within the husband's family if both of them held HIV positive status. Females are often blamed for the death of her husband/male partner, even if the male had been diagnosed HIV positive earlier. Females reuse needles and injecting equipment with their male partners," she added.

In the socio-cultural domain, drug use in Mizoram existed since long and casual sex has never been a taboo in some tribal culture. Many of the women drug users grew up in single parent households, experiencing an unstable and unhappy childhood [10]. Champai's geographical location reportedly gave rise to drug trafficking and human trafficking with Chins tribe people of Myanmar, stereotyped

as drug peddlers and traffickers, being and residing permanently in Champai district mostly [10]. The strong sense of community among the people in the North-eastern states has its negative aspects: drug users, people living with HIV/AIDS and especially female drug users, and their immediate families are often the target of force discrimination [10]. The numerous ethnic communities are affiliated to social and religious organizations such as Young Miso Association and Churches of northeast; but these institutions plagued by incomplete and incorrect information on drug use and HIV and holding narrow moral angle, too acted as a major source of stigma and discrimination. The CSC functionary stated, "Easy availability of illegal heroin pushed young people to opioid use and also the sudden surge in wealth and property, which is attributed to profits made from smuggling the precursors, and stress arising out lack of suitable employment opportunities educated youth in the region were contributing factors". Also, he pointed out, "Many of females into drugs also act as conduit for drug peddling and peer influence push them into IV drug use". An ARTC service provider stated "for FIDUs, treatment options through drug de-addiction and due coverage under targeted intervention programmes not existed. FIDUs are reluctant to go to oral substitution clinics attended by men; hence many female users stay out of reach of treatment facilities". According to CSC functionary "gender bias against FIDUs and more so against PLHIV FIDUs in general healthcare settings, barring ART centre, in Champai often compel FIDUs, requiring specialized treatment for certain ailments, suffer

Drug possession and peddling in Mizoram, is punishable offence and repeated case of Police arrests reported. Stringent law enforcement against heroin trafficking and peddling, in the early 1990s, resulted in shift towards the vein puncturing habit of injecting heroin and other pharmaceutical products. In the perspectives of the above drivers and facilitators, stigma 'marking', intersecting with socially discredited behaviour/characteristics on one side and HIV positive status identification and association with persons living with HIV (PLHIV) on the other hand was evident. This caused the emerging of stigma 'manifestation' for PLHIV individual and groups; and FIDUs carry the multiple layers of stigmatization i.e. gender norms, injecting drug use and HIV infection. The study thus found several forms of stigma, as stigma of being a drug user, gender-related stigma of being female injecting drug user, and stigma of being PLHIV.

In the perspectives of the above it is important to address different forms of stigma to mitigate their negative impacts on women's ability to access health services at individual, social and structural levels. At

imparted [16]. In addition, peer navigation to health facilities would reduce experiences of stigma and discrimination among FIDUs. The current harm reduction programme, supported by NACO across states, required to be refocused with inclusion of effective community and socially oriented harm reduction interventions in mitigating stigma, as it provides an avenue to strengthen employment, livelihood and skills development [16], progressive policing [17], legal support and violence mitigation with a rights-based approach [18,19,20].

In the current study, we embarked at understanding the stigma and discrimination issues concerning female injecting drug users based on the conceptualized stigma framework. This framework facilitated in analysing the social and structural pathways in addition to individual pathways. We observed that the stigmatization process unfolds across the socio-ecological spectrum and varies across economic contexts in low-, middle- and high-income countries [20]. We attempted to study the drivers, facilitators, intersecting stigma and stigma manifestations and observed that drivers and facilitators determine stigma marking through which stigma is applied to people or groups. We posited that stigma manifestations influence a number of outcomes for affected individuals and groups, including social acceptance, access to and uptakes of healthcare services, resilience and advocacy. The use of this framework enabled us to gauge stigma among PLHIV FIDUs in concise and comparable manner; and hopefully may be found suitable for use in future research studies, as well.

Limitations and Future Direction

Our study had certain limitations as it involved participants though registered under antiretroviral therapy regimen, but not availing harm reduction service coverage. They were linked to a care and support organization as such their accounts and experiences of stigma may differ from other women. It is indeed possible that our study may be underestimating the impact of stigma among female injecting drug users of Champai, because our purposively sampled participants were already accessing some psychosocial support and anti-retroviral treatment services.

Conclusions

Without intending to totally understand or cause oversimplification of the context and experiences of stigma, it is evident that women who injects drugs in Champai or anywhere in Mizoram often self-stigmatize, face stigma of injecting drug use, and are discriminated.

These are deterrent in their accessing and availing of health services. Suggestively, to overcome the multiple forms of stigma simultaneously experienced by participants in this study and ensure that tailored gender-sensitive interventions are available to them, a range of specific individual-, social-, and structural-level interventions will need to be implemented. For these to auger, necessary review of National Policy Narcotic Drugs and Psychotropic Substances 2012, to enable suitable inter-ministerial coordination on provision of addiction treatment under the Union Ministry of Social Justice, as well as access to Harm Reduction interventions and sexual health under Union Ministry of Health and Family Welfare are equally meted to female injecting drug users with suitable facilitation at state levels.

Acknowledgment

The study team acknowledges with thanks the support received from Participants and Healthcare Service Providers of Champai for their spontaneous and meaningful interactions. The authors are grateful for the help received from the Mizoram State AIDS Control

Society, Champai District AIDS Prevention Control Unit, New Hope Society, and Champai for facilitating the study.

Copyright: The manuscript has not been published in any Journal either a Research Paper and/or Conference Paper presentation. Authors do hereby grant Romanian Journal of Psychological Studies licensee to publish their work should it be considered for the same through the review process. Submission of a manuscript implies that the work described has not except in the form of an abstract or as part of a published lecture, been published before (or thesis) and it is not under consideration for publication elsewhere; that when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclosure statement

Ethical approval- The study had the approval of Mizoram State AIDS Control Society authority vide No. 11019/1/11/CMO (CPI)/DAPCU 2652 Dated 23.09.2019

Conflict of interest – Authors declare no conflict of interest.

Funding- This research received no specific grant from any funding agency, commercial entity or not-for-profit organization.

1. Úcæ } *|ÁŠkE } ^ÉÁÓæ! } •@æ , ÁÁKÉÁXæ| ^i^ÉÁŠ [*i^ÁPÁÓæ! { ^ }ÉÁÓiæ\ ^|Áçæ } Á Y i { ÉÁ^cæ|ÉÁ (2019) The health stigma and discrimination framework: a global, crosscutting, framework to inform research, and policy on health-related stigma, BMC Med 17:31
2. Silverman D (2001)