Short Communication Open Acces

Keywords: Carcinogenic e ect; Arsenate; P. vittata; Aqua-glycoporins

Metalloid arsenic has several structural form (Table 1), and can combine with many metals such as iron and molybdenum but Arsenic III (inorganic form) is most toxic form and is accumulating and entering in food chain constantly, escalating arsenic mobility issue more than calculated one. Further environmental condition such as pH, oods (redox condition), are conducive in high mobility and interconversions of arsenic getting complexed with many other compounds, favouring their easy transport in rice and other seeds via multiple transporters e.g., aquaglyceroporins in both plant and animals [1].

Marine organism had exceeding high levels of arsenic accumulation in non-toxic organic form arsenobetain, while rice is reported to have exceeding levels of inorganic arsenic (III). Arsenate As (V) is non-toxic form but it becomes toxic when arsenic (V) combines with phosphate or iron oxide and thus inhibits phosphorylation processes a er entering the cells. Arsenate As (III) causes deactivation of enzyme due to its high a nity towards thiol groups. In addition, inorganic arsenic enters the cell via the hexose transporter, phosphate transporter systems (PTS) or aqua-glycoporins, while rice and other plants had di erent transporter. Arsenic reductase coded by Ars or Arr operon are helpMC /l-6(ru-5(l vn no)12(mn)23v)8(e)s on of ersenic12.1(n)6(e t)-61(o)arsenote tn nb-8.9(o)1109(t)66(h)rokdsc

Citation: Bhatt SM (2016) Arsenic as Next Global Threat? Role of Biotechnological Approaches. J Bioremed Biodeg 7: 329. doi: 10.4172/2155-6199.1000329

© ½ º Ã » ³ Ú œ ÁÃ » 0 0 • 0

40.

© ½ º Ã » ³ Ú œ ÁÃ » 0 0 • 0