Vromman et al., J Bioremediat Biodegrad 2016, 7:3
DOI: 10.4172/2155-6199.1000349

Research Article Open Access

***Corresponding author:** Stanley Lutts, Groupe de Recherche en Physiologie

e site is located at the border between the rst and second Region of Chile. In the environment of *Atriplex atacamensis*, trees such as *Prosopis sp*., *Geoffroea decorticans* and aquatic plant as *Scirpus americanus*, *Ruppia maritime* and *Stygeoclonium sp.* were also observed.

Fourteen plants of *A. atacamensis* were identi ed (10 female and 4 male plants) on a surface area of 0.25 ha. e soil samples (approximately 500 g each) were taken to a depth of 0 to 40 cm (4 st7e s (tho7o obeEMC /Span <</Lang (en-US)/MCID 270 ЭDC 1BT0.034 Tw 9 0 0 9 42.5197 710.568

WHO for drinking water. e water is also highly saline, with a mean EC value higher than 14 mS cm^{-1} in relation to a high concentration of Na and Cl. Similar As values in the Rio Loa were reported by Bugueño et al. [9] and Romero et al. [2]. According to these authors, the As concentration in water of Rio Loa is quite lower in the upper section and progressively increased from the Andes mountain to the paci c ocean, as a consequence of extreme desert climate favoring high evaporation rates and the lack of tributaries or fresh groundwater contribution. Such As-enrichment thus occurs as a natural process, although some anthropogenic activities related to mining ore processing may also be involved. Beside As, water also contains high amounts of B, and a near-neutral to alkaline pH. ose pH values, high salinity and high As concentrations do not favour the adsorption of As(V) species [22]. Considering the mean redox potential, both $HAsO_4^-$ and H_3AsO_3 may be present in the water [23]. At pH<7 in oxidizing conditions, As may be immobilized by co-precipitation/sorption with Fe and Mn oxy-hydroxides [24]. Despite a high total dissolved solid concentration (TDS), Fe and Mn are poorly represented in the water of Rio Loa. Similarly, under acidic sulfate reducing conditions, As can be accumulated as stable sulfur minerals [25] which probably does not occur at the pH recorded in the samples. For Ca/As molar ratio higher than 1 and low dissolved Fe and Mn concentration, As may be associated with calcite and kaolinite minerals [26] and the presence of this complex in suspension could not be ruled out under the recorded conditions.

Data issued from bulk soil analysis are given in Table 2. Soils of Quillagua were slightly alkaline and highly saline, containing only a small amount of organic matter. ey were typically coarse-textured, dominated by medium and ne sand. e total concentration of As was high (220 mg Kg^{-1}), which is by far higher than the world average concentration reported for this matrix [27]. Higher vEMC /Spay9,n conlng (eg)8 (e)]10 (t)6 (e)ion c[q ƀBDC B6was higrde

Figure 3: A)

by high As concentration in the soil, but also by high B availability, high salinity, high temperature and very high vapour pressure de cit. is species thus appears resistant to as a promising candidate for the phytomanagement of contaminated soils under arid conditions.

References

1. Queirolo F, Stegen S, Mondaca J, Cortés R, Rojas R, et al. (2000) Total arsenic, lead, cadmium, copper, and zinc in some salt rivers in the northern Andes of Antofagasta, Chile. Sci Total Environ 255: 85-95.
2. Romero L, Alonso H, Campano P, Fanfani L (2003) Arsenic enrichment in waters and sediments of the Rio Loa (Second Region, Chile). Appl Geochem 18: 1399-1416.
3. Díaz O, Tapia Y, Pastene R, Montes S, Núñez N, et al. (2011) Total and bioavailable arsenic concentration in arid soils and its uptake by native plants from the pre-Andean zones in Chile. Bull Environ Contam Toxicol 86: 666-669.
4. Karcher S, Cacéres L, Jerkel M, Contreras R (2007) Arsenic removal from water supliesq in Northern Chile using ferric chloride coagulation. J Chart Inst Water Environ Manag 13: 164-169.
5. Pell A, Márquez A, Rubio R, López-Sánchez JF (2013) Effect of sample processing on arsenic speciation in marine macroalgea. Anal Meth 5: 2543-2550.
6. Smedley PL, Kinniburgh DG (2013) Arsenic in Groundwater and the Environment. British Geological Survey.
7. Smith AH, Goycolea M, Haque R, Biggs ML (1998) Marked increase in bladder and lung cancer mortality in a region of Northern Chile due to arsenic in drinking water. Am J Epidemiol 147: 660-669.
8. Smedley PL, Nicolli HB, Macdonald DMJ, Barros AJ (2002) Hydrogeochemistry of arsenic and other inorganic constituents in groundwater from La Pampa, Argentina. Appl Geochem 17: 259-284.
9. Bugueño MP, Acevedo SE, Bonilla CA, Pizarro GE, Pasten PA (2014) Differential arsenic binding in the sediments of two sites in Chile's lower Loa River basin. Sci Total Environ 466-467: 387-96.
10. Tripathi RD, Srivastava S, Mishra S, Singh N, Tuli R, et al. (2007) Arsenic hazards: strategies for tolerance and remediation by plants. Trends Biotech 25: 158-165.