

" TTFTTNFOUT PG )FBWZ .FUBM "DDVNVMBUJPO  
4QFDJFT GPS 1IZUPSFNFEJBUJPO " \$BTF 4UVE

Mekonnen Amberber <sup>1</sup> DQG )DQWX .LÀH

<sup>1</sup>Department of Environmental Science, Kotebe University College, Addis Ababa, Ethiopia

<sup>2</sup>Addis Ababa Environmental Protection Authority, Addis Ababa, Ethiopia

## Abstract

that Cr (153, 22 mg/L), Ni (30.71 mg/L) and Pb (9.55 mg/L) were detected from the soil but Cd was not detected. The plant analysis result indicated that the species *Arundo donax*, *Ricinus communis* and *Vernonia amygdalina* had a good potential plant that absorbs Cd, Cr and Pb in different concentrations, while Ni was not detected in the leaves of selected plants. The species *Arundo donax* accumulated Cr (80.90 mg/L), Pb (37.30 mg/L) and Cd (25.98 mg/L), *Vernonia amygdalina* accumulated Cr (83.59 mg/L), Cd (44.46 mg/L) and Pb (14.49 mg/L) and *Ricinus communis* accumulated Cr (62.06 mg/L), Cd (16.64 mg/L) and Pb (16.64 mg/L). It was concluded that the selected plant species had a good accumulation capacity of Cr, Pb and Cd for the phytoremediation activity.

**Keywords:** Accumulation capacity; Atomic absorption nickel using agricultural crops including ricinus using three levels of spectrophotometer; Heavy metals; Little Akaki River; Phytoremediation

## Introduction

Consequent to global industrialization, heavy metal pollution is a widespread problem which has become a major environmental concern due to hazardous effects on human and environmental health [1,2]. Air and water pollution by metals varies from soil pollution, because heavy metals persevere in soil for a longer time period as compared with the other compartment of the biosphere [3]. In the latest decades, the yearly global release of heavy metals attained 22,000 t (metric ton) for cadmium, 939,000 t for copper, 783,000 t for lead, and 1,350,000 t for zinc [4-12].

Many phytoremediation technologies have been used for the remediation of polluted soils and water throughout the World [1,5]. Phytoremediation costs almost one-fourth of the other physical and chemical methods of pollutant treatment [5]. The major advantages of the process include: improvement of the soil quality, as it is driven by solar energy thus suitable to most regions and climates, cost effective and technically feasible process, plants serve as sufficient biomass for rapid remediation; promote high rhizosphere activity and finally restoration in a reasonable time frame of 2 to 3 years [1,2].

Some plants, which are often identified as bioaccumulators, have the ability to take up soil contaminants and deposit them in their roots, as well as in their aboveground organs. According to Kowalska [6,10] it is necessary to point out that bioaccumulating plant species are normally characterized by high concentration factors, i.e., concentrations of the toxic substances are higher in their tissues than in the soil. Bioaccumulation factors of some plants can even reach 1000x. Some plants are capable of intensive uptake of soil contaminants and, at the same time, are characterized by a significant production of biomass [6].

Previous experiments on giant reed suggested that the stem height and diameter, number of nodes, fresh and dry weight of leaves, and net photosynthesis were not affected, indicating that plants tolerated the high concentrations of Cd and Ni [7,9]. On the other hand, a study conducted by Ref. [6,7]. Phytoremediation of soil polluted by

Gulele sub city and Nefas silk la o sub city respectively, Addis Ababa, Ethiopia. The four sample sites were selected based on preliminary survey (physical observation and discussion with woreda and sub city expert) and literature review than delineated as illustrated (Table 1 and Figure 1).

Corresponding author: Mekonnen Amberber, Department of Environmental Science, Kotebe University College, Addis Ababa, Ethiopia, Tel: +251913714680; E-mail: [mekonnenamberber@yahoo.com](mailto:mekonnenamberber@yahoo.com)

Received July 13, 2016; Accepted August 26, 2016; Published August 27, 2016

Citation: Amberber M, et al. (2016) Assessments of Heavy Metal Accumulation Capacity of Selected Plant Species for Phytoremediation: A Case Study in Little Akaki River. *J Bioremediat Biodegrad* 7: 369. doi: [10.4172/2155-6199.1000369](https://doi.org/10.4172/2155-6199.1000369)

Copyright: © 2016 Amberber M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

pollutant discharging. Soil samples were randomly collected within the river bank whilst plant leaf samples were collected from dominant plant that was growing along the river banks [4,8,9].

#### Data analysis methods

The elements of interest for these particular analyses were four (4) heavy metals and they are cadmium (Cd), lead (Pb), Chromium (Cr) and Nickel (Ni). All samples were analyzed using Atomic Absorption

is experimental result indicated that Ni was not detected in all heavy metals (Pb, Cr, and Cd) were detected in the tested plant leaves samples of leaf. This finding agrees with the result of Yanqun et al. [7] in different concentration, whereas Ni was not detected. According to and Alebachew [13] the possible reasons are listed by him is that Ni may be degraded in soil by microorganism activity (Phytodegradation) or stored other part of the plants (phytostabilization) or both of them. Hahun [12] the possible reasons to this concentration difference are growth rate of plants, their ability of absorbed, accumulated capacity of heavy metals in their parts and the depth of root zones or other. As illustrated in Figure 2, *A. ... a a ... a ... a* had good potential of absorbing and accumulating Cr than the others. While *... a* had relatively good capacity to store Pb than the others.

## Conclusion

This study generally showed that the leaves sample of *A. donax* and *A. donax* had high potential of absorbing Cr and Pb comparatively. Whereas the leaves of *A. donax* had high potential of absorbing Cr and Pb paralleled to others plant species. Cd, Pb, Ni and Cr was detected in all sampled (soil, water and leaves), whereas Ni was not detected from all sampled plant species. Therefore planting those plants around polluted river bank for the remediation of Cr, Pb and Cd is very important. This study was conducted in field leaf samples, an extended and detailed experimental study in a controlled manner is necessary.

## References

1. Yanqun Z, Yuan L, Schwartz Ch, Langlade L, Fan L (2004) Accumulation of Pb, Cd, Cu and Zn in plants and hyperaccumulator choice in Lanping lead-zinc mine area. *China Environ Int* 30: 567-576.
2. Gisbert G, Ros R, Haro AD, Walker DJ, Bernal MP, et al. (2003) A plant JHQHWLFDQO\ PRGL¿HG WKDW DFFXPXODWHV 3E LV HVSHFLDOO\ SURPLVLQJ IRU phytoremediation. *Biochem Bioph Res Co* 303: 440-445.
3. Lasat MM (2002) Phytoextraction of toxic metals: A review of biological mechanisms. *J Environ Qual* 31: 109-120.
4. Singh OV, Labana S, Pandey G, Budhiraja R, Jain RK (2003) Phytoremediation: an overview of metallic ion decontamination from soil. *Appl Microbiol Biotechnol* 61: 405-412.
5. DPUDQ O\$ \$PQD OXIWLHWDO 5 7KH SRWHQWLDO RI WKH ÀRUD IURP GLIIHUHQW regions of Pakistan in phytoremediation: a review. *Environ Sci Pollut Res* 21: 801.
6. Álvarez E, Fernández Marcos ML, Vaamonde C, Fernándezsanjurjo MJ (2003) Heavy metals in the dump of an abandoned mine in Galicia (NW Spain) and in the spontaneously occurring vegetation. *Sci Total Environ* 313: 185-197.
7. Papazoglou EG, Karantounias GA, Vemmos SN, Bouranis DL (2005) Photosynthesis and growth responses of giant reed (*Arundo donax* L.) To the heavy metals Cd and Ni. *Environ Int* 31: 243-249.