manifestations. Some of the most disruptive behaviors, referred to as aberrant behaviors, can cause signiýcant disability and distress to the patient and caregiver (Baghdadli, Pry, Michelon, & Rattaz, 2014). Aberrant behavior is divided into subcategories by the Aberrant Behavior Checklist (Slosson Educational Publications Inc, East Aurora, NY). These categories include Irritability, Social Withdrawal, Stereotypy, Hyperactivity and Inappropriate Speech. Irritability, which includes severe tantrums, aggression, and self-injury, is one of the major and most disruptive aberrant behaviors (Stigler, 2014). Irritability is commonly treated with antipsychotic medication with or without behavior therapy (Aman et al., 2009). Treating with a medication to suppression symptoms rather than understanding if there is an underlying medical cause for such behaviors can be problematic for several reasons. First, treating symptoms instead of identifying a cause can lead to increased morbidity and mortality from an undiagnosed disorder. Second, antipsychotic medications can detrimentally affect glucose, cholesterol and lipids and weight, even in the short-term (Correll et al., 2009; Wink et al., 2014) and long-term anti-psychotic use increases the risk for type II diabetes (Bobo et al., 2013) and can result in tardive dyskinesia, a potentially permanent movement disorder (Correll & Kane, 2007). Thus, it