




---

**\*Corresponding author:** Antonio Eduardo De Aquino Junior, São Carlos Institute of Physics, University of São Paulo, São Carlos, São Paulo, Brazil, Tel: +55-16991721127; E-mail: [antonioaquino@ifsc.usp.br](mailto:antonioaquino@ifsc.usp.br)

**Received** May 03, 2020; **Accepted** May 21, 2020; **Published** May 28, 2020


**Citation:** Tamae PE, Santos AVD, Simão MLS, Canelada ACN, Zampieri KR, et al. (2020) Can the Associated Use of Negative Pressure and Laser Therapy Be A New

## Fundamentals for a Therapeutical Approach Involving Negative Pressure and Laser




---

<b>Therapy</b>	<b>Parameters</b>
<b>Laser Therapy Group</b>	Application sites consist of the paravertebral region, extending from the cervical spine to the lumbar spine and scapular waist bilaterally in the dorsal trunk, with the application being scanned, 5 min in the right region and left region of the cervical region and dorsal trunk. In the upper limbs, in an anatomical position, the application sites were 5 min on the arm



## Discussion

The text in this section is also illegible due to the same quality issues as the figure above. It appears to be a dense block of text, but the characters are too small and blurry to read.



**Citation:** Tamae PE, Santos AVD, Simão MLS, Canelada ACN, Zampieri KR, et al. (2020) Can the Associated Use of Negative Pressure and Laser Therapy Be A New and Efficient Treatment for Parkinson's Pain? A Comparative Study. *J Alzheimers Dis Parkinsonism* 10: 488.

---