



**Keywords:** Impaired fasting glucose; Impaired glucose tolerance; Prediabetes complications; Cardiovascular Disease; Macro vascular

## Introduction

According to the World Health Organization (WHO), one in ten adults worldwide had type 2 diabetes mellitus, which is one of the main causes of premature morbidity and mortality. Diabetes-related healthcare accounts for one out of every five dollars spent on health care in the United States [1]. Additionally, macro vascular complications account for the majority of the direct and indirect costs associated with diabetes in developing nations, accounting for at least a third of their total expenditures on diabetes-related health care. An interlude of prediabetes, itself a toxic state that is linked to the development of macro vascular complications, occurs during the T2DM progression [2].

## Method

### Key causes and effects of prediabetes

Impaired glucose tolerance (IGT) is defined by the American Diabetes Association as a 2-hour plasma glucose value in the 75-gram oral glucose tolerance test of A fasting plasma glucose level between 100 and 125 mg/dL is considered impaired fasting glucose (IFG).6 Finally, prediabetes can also be characterized by a haemoglobin A1c level between 5.7% and 6.4%. It is important to point out that the ADA criteria define normal glucose tolerance (NGT) as having a fasting

and adiponectin levels in the blood as well as higher triglyceride levels.

### Cardiovascular disease

In the EPIC-Norfolk study, a higher 10-year cardiovascular mortality rate was linked to an increase in within the normal range. The EPIC-Norfolk results are consistent with the Paris Prospective Study cohort data, which showed that IGT subjects had a doubling of CVD mortality compared to NGT subjects [7]. The majority of patients with prediabetes have features of insulin resistance (metabolic) syndrome, such as upper-body obesity, hypertriglyceridemia, decreased HDL cholesterol levels, and hypertension, among other things. This highlights the finding of increased mortality. In prediabetes patients, components of the metabolic syndrome can frequently be identified years before a diagnosis is made. These characteristics result in advanced atherosclerotic vascular changes, which are frequently preceded by impairment of endothelium-dependent vasodilation, dysfunction of vascular smooth muscle, and elevated arterial stiffness [8]. A recent cross-sectional study found a positive correlation between the prevalence of arterial stiffness and prediabetes, indicating that prediabetes control may prevent arterial stiffness with early intervention. According to the Multi-Ethnic Study of Atherosclerosis (MESA) study, the prevalence of unrecognized myocardial infarction is almost three times higher in people with prediabetes than in people with NGT. In a more recent randomized controlled trial involving 6522 people with prediabetes and coronary artery disease, it was discovered that although acarbose did not lower the risk of major adverse cardiovascular events, it did lower the incidence of diabetes.

### Conclusion

Patients should be routinely tested for prediabetes by doctors and other healthcare professionals, and those who have the condition should be referred for intensive lifestyle counselling. Similar to the DPP and Kindred studies, the objective is to lose more than 5% of body weight and keep it off through calorie restriction and increased physical activity [9]. In order to increase the availability of structured lifestyle programs that are based on evidence, healthcare providers ought to make an effort to establish strong ties within healthcare systems, communities, and payers. Based on HbA1c results, a recent survey of outpatients found that many had prediabetes. Amazingly, clinicians recognized and diagnosed prediabetes in those patients whose HbA1c tests showed it. Only 23% of those with prediabetes

whose status was accurately documented in the clinical records had treatment documentation [10]. During the prediabetes phase, it is possible to intervene in the disease process. Patients, specialists, health systems, and primary care physicians all have a responsibility to prevent missed opportunities for prevention. Although it has long been known that diabetes poses significant risks to the cardiovascular system, it is now becoming clear that people with prediabetes are more at risk for CVD than those with diabetes. The impact on cardiovascular health is staggering when considering the millions of prediabetes individuals worldwide. However, education, increased awareness, care coordination, organization, and novel reimbursement mechanisms at multiple levels are necessary for identifying and intervening in the at-risk populations of prediabetes.

1. Dall TM, Yang W, Halder P (2014) The Economic Burden of Elevated Blood Glucose Levels in 2012. *Diabetes Care* 37 (12): 3172–3179.
2. Dagogo-Jack S (2010) Pitfalls in the use of HbA1(c) as a diagnostic test: the ethnic conundrum. *Nat Rev Endocrinol* 6(10): 589–593.
3. Chapp-Jumbo E, Edeoga C, Wan J, Dagogo-Jack S (2012) Ethnic disparity in hemoglobin A1C levels among normoglycemia of spring of parents with type 2 diabetes. *Endocr Pract* 18(3):356–362.
4. Ebenibo S, Edeoga C, Wan J, Dagogo-Jack S (2014) Glucoregulatory function among African Americans and European Americans with normal or pre-diabetic hemoglobin A1C levels. *Metabolism* 63(6): 767–812.
5. Soranzo N (2011) Genetic determinants of variability in glyated hemoglobin (HbA (1c) in humans: review of recent progress and prospects for use in diabetes care. *Curr Diab Rep Dec* 11(6): 562–600.
6. Herman WH (2016) Are There Clinical Implications of Racial Differences in HbA1c? Yes, to Not Consider Can Do Great Harm! *Diabetes Care*. Aug 39(8):1458–1505.
7. Middelbeek RJW, Abrahamson MJ (2014) Diabetes prediabetes and glycaemic control in the United States: challenges and opportunities. *Ann Intern Med* 160 (8):572–573.
8. Hu D, Fu P, Xie J (2008) MS for the InterASIA Collaborative Group Increasing prevalence and low awareness, treatment and control of diabetes mellitus among Chinese adults: the InterASIA study. *Diabetes Res Clin Pract* 81(2):250–257.
9. Saadi H, Carruthers SG, Nagelkerke N (2007) Prevalence of diabetes mellitus and its complications in a population-based sample in Al Ain, United Arab Emirates. *Diabetes Res Clin Pract* 78 (3)369–377.
10. Echouffo-Tcheugui JB, Dagogo-Jack S (2012). Preventing diabetes mellitus in developing countries. *Nat Rev Endocrinol* 8 (9):557–562.