Central Node Neck Dissection for Papillary hyroid Carcinoma: Clinical Implications, Surgical Complications and Follow up. A Prospective vs a Restrospective Study

Maria Rosa Pelizzo[1], Isabella Merante Boschin[1*], Maddalena Variolo[1], Giovanni Carrozzo[1], Costantino Pagetta[1], Ornella Lora[2], Andrea Piotto[1] and Chiara Dobrinja[3]

[1]II Surgical Clinic Unit, Department of Oncological Surgical and Gastroenterological Sciences (DiSCOG), University of Padova, Via Giustiniani 2, Padova, Italy

[2]Radiotherapy and Nuclear Medicine Unit, Istituto Oncologico Veneto-IRCCS, Padova, Italy

[3]Department of Medical and Surgical Sciences, Division of General Surgery, Hospital of Cattinara, University of Trieste, Italy

[*]**Corresponding author:** Isabella Merante Boschin, Unit, Department of Oncological Surgical and Gastroenterological Sciences (DiSCOG), University of Padova, Italy Tel: +0390498212250; E-mail: isabella.meranteboschin@unipd.it

Received date: July 21, 2015; **Accepted date:** December 01, 2015; **Published date:** December 03, 2015

Copyright: © 2015, Pelizzo MR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: The treatment and particularly the extension of surgical therapy of papillary thyroid carcinoma (PTC) remain still controversial in some issues, especially for the lack of preoperative information or variables that allow predicting the level of aggressiveness of the tumor.

Aim of the study: The purpose of the study was to assess the impact of the central node neck dissection (CNND) on surgical outcome and disease free- follow up of PTC- patients operated on at our center by evaluation of postoperative complications (parathyroid and recurrent nerve damage, hemorrhage rates) and pts rates presenting detectable serum Thyroglobulin (TG) or TG-Antibodies (TG-AB) values, at the time of ^{131}Iodine treatment and
'the
ae