

Changes in the Frequency of Airway Infections in Smokers Who Switched To Vaping: Results of an Online Survey

Joanna Astrid Miler^{1*}, Bernhard Mayer² and Peter Hajek¹

¹Wolfson Institute of Preventive Medicine, Queen Mary University of London, London, UK; ²Department of Health and Lifestyle Research, Queen Mary University of London, London, UK; *Corresponding author: j.miler@qmul.ac.uk

Received date: July 04, 2016; **Accepted date:** July 30, 2016; **Published date:** Aug 05, 2016

Copyright: © 2016 Miler JA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background and aim: Cell and animal studies suggested that use of e-cigarettes may increase vulnerability to respiratory infection, though the available studies have serious limitations. Limited data are available on respiratory health of vapers.

Methods: An on-line survey assessed subjective changes in respiratory symptoms in smokers who switched to vaping for at least two months.

Results: Among 941 responders, 29% reported no change in respiratory symptoms, 5% reported worsening, and 66% reported an improvement. Among qualitative comments, 232 elaborated on positive and 15 on negative experiences.

Conclusion: The switch from smoking to vaping was associated with a reduced incidence of self-reported respiratory infections. Further studies using objective measures in samples that are not self-selected are needed.

Keywords: E-cigarettes; Vaping; Airway infections; Smoking cessation; Survey

Introduction

E-cigarettes (EC) deliver nicotine without combustion chemicals and are therefore considered to be much safer than conventional cigarettes [1,2]. Some potential risks however have been identified in cell and animal studies suggesting that vaping may increase vulnerability to respiratory infections [3,4]. There are only limited data available on respiratory health of vapers. Human EC trials reported no adverse respiratory symptoms associated with EC use for up to 1.5 years and a follow-up study of smokers with asthma who switched to vaping found improvements. We conducted an on-line survey to assess subjective changes in respiratory symptoms in smokers who switched to vaping for at least two months [5,6].

Methods

Design

The survey was posted by BM on two large German vape forums previously used by BM, E-Rauchen-Forum (<http://www.e-rauchen-forum.de/thread-82289.html>) and E-Zigaretten-Forum (<http://www.e-zigaretten-forum.de/thread-10000.html>). Most of the active members of these forums are ex-smokers using tank system EC products.

The pre-ambles to the survey explained that a recent article suggested that vaping could increase rates of common cold and respiratory

infections; that BM's experience was the opposite; and that the purpose of the survey is to find out about any changes in rates of respiratory infections vapers experience [3]. Responders were asked to respond honestly and to note that the survey concerns respiratory infections such as common cold and no other respiratory diseases such as asthma or COPD. Responses were invited only from vapers who stopped smoking completely or reduced smoking by at least 95% and who have been vaping for at least 2 months. Appendix A includes verbatim translation of the pre-ambles and a screenshot of the survey page.

The question asked was: Since switching to e-cigarettes, the frequency of any airways infections I get has: a) decreased; b) not markedly changed; c) increased. We calculated proportions for each answer and their 95% confidence intervals for each answer using SPSS.



Figure 1: Self-reported changes in airway infections U Yf switching from smoking to vaping (N=941).

YfY were 247 qualitative responses commenting on improvements or worsening in the rates of respiratory infections U Yf switching from smoking to vaping Y total of 232 comments concerned positive and 15 concerned negative Y YMg

Here are some typical reports of positive changes: “As a smoker I had an almost continuous cold. I’ve been a vaper for almost 2 years, and for almost 2 years I’ve had no gb] b[žno cough, nothing..”

“I’ve been vaping for 13 months and had none [infections] while earlier I had colds and sore throat 2-3 times a year.”

“I’ve been vaping for 2 years and in this time I had one i !]_Y infection. As a smoker I used to have them at least twice a year, for weeks at a time. I used to have to deal with a persistent cough and mucus production.]gyear the infection had passed within 3 days.”

Here are some examples of negative reports:

“I smoked for 15 years and in those 15 years I was never really ill. 5 Yf quitting smoking I have already had 2 i !]_Y infections [...].”

“I have completely switched to vaping in 2009. At that time, the fgh 3 smoke-free years I was XY b]l]Ymsick less c Yb than during my smoking years. But I have observed that I have been sick more c Yb in the last 3-4 years (a cold about 4-5 times a year). However, I must say that I became a father 4 years ago. Maybe it has to do with my daughter - she c Yb passes on her infections to me. I’m mostly ill U Yf she has been ill.”

Discussion

Y

- 2 Tobacco Advisory Group of the Royal College of Physicians (2016) Nicotine without smoke-tobacco harm reduction. Royal College of Physicians
- 3 Wu Q, Jiang DJ, Minor M, Chu HW (2014) Electronic cigarette liquid increases *Staphylococcus aureus* and virus infection in primary human airway epithelial cells. *PLoS One* e108342
- 4 Sussan TE, Gajghate S, Jaisankar UR, Ma J, Kim JH, et al. (2015) Exposure to electronic cigarettes impairs pulmonary anti-bacterial and anti-viral defenses in a mouse model. *PLoS One* 10: e0116861.
- 5 McRobbie H, Bullen C, Hartmann-Boyce J, Hajek P (2014) Electronic cigarettes for smoking cessation and reduction. *Cochrane Database Syst Rev* CD010216
- 6 Polosa R, Morjaria J, Caponnetto P, Caruso M, Strano S, et al. (2014) Effect of Smoking Abstinence and Reduction in Asthmatic Smokers Switching to Electronic Cigarettes: Evidence for Harm Reversal. *Int J Environ Res Public Health* 11: 4965-4977.
- 7 Feldman C, Anderson R (2013) Cigarette smoking and mechanisms of susceptibility to infections of the respiratory tract and other organ systems. *J Infect* 67: 169-184
- 8 Nalawade TM, Bhat K, Sogi SH (2015) Bactericidal activity of propylene glycol, glycerine, polyethylene glycol 400 and polyethylene glycol 1000

1 D V P À` `