

Characteristic of Neuronspecific Markers in Preterm Infants with Hypoxic-ischemic Encephalopathy

Department of Neurology, Yerevan State Medical University, Yerevan, Armenia

* : Lilit Galstyan, Department of Neurology, Yerevan State Medical University, Yerevan, Armenia, Tel: 37410484280; E-mail: [Ugnib`]l4na U`Wa

: February 05, 2018; : February 26, 2018; : March 07, 2018

: © 2018 Galstyan L. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Problems of diagnosis of hypoxic-ischemic encephalopathy in preterm neonates remain relevant, serious and not fully studied and not fully refined directions.

The revealed regularities and peculiarities depending on the gestational age of newborns allows as to assert that applied immune-fermentative studies for the diagnosis of hypoxic-ischemic encephalopathy show that they need further improvement, and instrumental methods are not allowed to have an early and clear understanding about the prevalence of pathological process, predict outcome and to develop therapeutic tactics.

< ncl JWgWá JW YbWd\UcdUhm' BY fcb'
a U_Yg'Dfj YbfJW'f~Y_ca UWU'7YbfU'bYj ci ggnVá

-i lg_bck b'hUhYbYk Vcfb'VU]Yg k]h' ck 'UbX]Yfm'ck 'VcXm
kY[\h a U Y i d' h Y a cgi \|]\!f]g 'dYf]bUH' XUhg UbX Wgg cZ
XgU]lm[fc i d'QQUa cb['h Y a 'h YdYf]bUH'a cfU]hriUYlg%!&
ha Yg \|]\!Yf hUb' h Y gla Y]bX]Wcf' cZ VU]Yg k]h' & SS' ['Vfh
kY[\h QQ

· gi XmcZ]HYUi fY XHU OQg\ckg hUi Zf' h Y a YbQck \

