23-Dec-2022, DOI: 10.4172/2332-0702.1000350

Methods

A comparative study was conducted using a consecutive sampling technique among 200 adult patients seeking medical and dental care. Data were collected from patients attending the Dental and Medical OPD of Dow University of Health Sciences in March 2015. Adult patients seeking health care without physical or mental disabilities aged between 20-70 years participated. Written informed consent was attained from each participant; permission to conduct the study was taken by the head of the hospital.

Data was gathered through Performa, which comprises demographic information and a valid oral health literacy instrument (OHL-I) [31]. One examiner collected demographic and oral health literacy information, and this process endures approximately 10 minutes. 200 patients participated in the study with a response rate of 99.4%.

e study inquired about the patient's gender, age, level of education, marital status, place of residence, and monthly income. e Oral Health Literacy Tool was translated into Urdu for a better understanding of patients. e oral health knowledge test used in the study included 10 statements related to dental caries, periodontal disease, oral cancer, and oral hygiene (Table 1). Oral health behavior was recorded as regularity of dental visits categorized as yes, don't remember, and never. Subjects were asked to rate each statement as true or false; correct answers were scored 1 and incorrect or don't know answers were scored 0. e total oral health knowledge score varies from 0 to 10 (the sum of scores for each item). e 10 statements that were recorded on Performa were entered into SPSS version 16. Oral health literacy was compared using an independent t-test, and chi-square was applied to categorical variables for the association between knowledge of a speci c question among patients.

Result

A total of 200 patients with a mean age of 32.8 ± 12.9 years, 59% women and 49% men were included. In terms of education, (35.5%) of the respondents graduated from elementary school, (20.5%) completed high school and professional graduation. (11.5%) ended middle school and only (5%) were postgraduate. Participants with a monthly income of less than PKR 10,000 were (17%), more than half of the participants have 10,000 monthly incomes (56.5%) and more than 10,000 were (26.5%) as shown in Table 2.

An independent samples t-test was used to compare oral health literacy among dental and medical patients. ere was a signi cant difference in oral health literacy scores for dental (M= 6.20, SD= 1.56) and medical (M= 5.43, SD=1.84) patients; t (198) = 3.18, p = .02 (Table 2). ese results suggest that oral health literacy di ers between dental and medical patients. Speci cally, the results indicate that dental patients when attending dental OPD their status of oral health literacy