Journal of
Obesity & Weight Loss Therapy

Choi, J Obes Weight Loss Ther 2015, 5:2
10.4172/2165-7904.1000255

Research Article Open Access

# Comparisons of 6-n-Propylthiouracil (PROP) Sensitivity, Food Liking and Food Intake between Vegetarian and Non-Vegetarian Women

**Sung Eun Choi***

*Department of Family, Nutrition and Exercise Sciences, Queens College, the City University of New York (CUNY), Flushing NY, USA*

***Corresponding author:** Sung Eun Choi, PhD, RD, Department of Family, Nutrition and Exercise Sciences, Queens College, CUNY Remsen Hall 305 A, 65-30 Kissena Boulevard, Flushing NY, 11367-1597, USA, Tel: 1-718-997-4169; Fax: 1-718-997-4163; E-mail: sungeun.choi@qc.cuny.edu

**Received date:** March 16, 2015; **Accepted date:** April 4, 2015; **Published date:** April 25, 2015

**Copyright:** © 2015 Choi SE. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Background:** Previous studies have suggested that vegetarians have a lower risk of overweight and obesity than do non-vegetarians. However, little is known about how meat consumption interacts with taste perception, thereby influencing food intake and body weight.

**Objective:** The objective of this study was to examine the relationship of meat consumption with 6-n-propylthiouracil (PROP) sensitivity, food liking, food intake and body mass index in female long term vegetarians and non-vegetarians.

**Methods:** A cross-sectional design with a total of 94 racially diverse female subjects (mean age 23 years, 42 vegetarians, 52 non-vegetarians) living in the New York City area was used in this study. Body Mass Index (BMI) was calculated using the measured weight and height and PROP sensitivity was assessed using a PROP filter paper disk method and a general Labeled Magnitude Scale (gLMS). The subjects also completed a questionnaire to report the food liking/disliking for 19 food items using a hedonic version of the gLMS. Dietary intake was assessed using a food frequency questionnaire.

**Results:** The PROP sensitivity of vegetarians was significantly higher than that of non-vegetarians. Vegetarians showed significantly less liking of fat foods than did non-vegetarians, whereas there was no significant difference in sweet foods liking. The BMI, energy, protein, fat, saturated fat, cholesterol, B vitamins, iron, zinc, sodium, potassium, and alcohol intake values of the vegetarians were significantly lower than those of the non-vegetarians. Multiple regression results revealed that only vegetarian status significantly contributed to the predictions of all the dependent variables, energy intake, fat intake and BMI.

**Conclusion:** The study findings suggest that a difference in taste perception between vegetarians and non-vegetarians may play a role in determining energy intake and weight status. Further studies are needed to examine the mechanisms by which habitual meat consumption affect taste perception and thus food intake.

**Keywords:** Vegetarian; Body mass index; Food liking, 6-n-propylthiouracil (PROP), Food intake

A vegetarian diet has been suggested as an approach for weight management by scientific literature. The scientific literature suggests that a plant-based diet is inversely related to Body Mass Index (BMI), and to the incidence of overweight, and obesity [1-3]. Although the studies in the reviews varied in their design and adjustment for potential confounders in the analyses, most studies found that vegetarians have a lower BMI on average [1-6]. Several studies observed that vegans have a lower mean BMI than that of other types of vegetarians [1,7-9]. A possible explanation for the obesity preventive effect of meatless eating patterns includes the consumption of lean protein foods, reduced-fat dairy products, dietary fiber and whole grains [2,10-13].

For weight-loss seekers to receive such potential benefits from a vegetarian diet, the subjects must strictly commit to a vegetarian diet for a significant period of time. However, studies showed that many vegetarians fail to strictly adhere to meat abstention [14-20]. Several large-scale surveys using a representative American sample also revealed that more than half of self-identified vegetarians admitted that they had eaten animal flesh [21-24]. In a study examining possible reasons for the discrepancy between self-definition and admitted behavior among vegetarians, Rothgerber suggested the weaker disgust for meat of non-strict vegetarians than that of strict vegetarians can be a cause or consequence of their occasional consumption of animal flesh [25]. Disgust is a negative emotion characterized by a primitive revulsion at the prospect of bodily contamination and a symbolic element rejecting immoral or polluting objects, behaviors, or individuals [26]. A study by Fessler identified disgust as a multifaceted emotional state that is potentially measurable as a mental state, contamination potency, nausea, ideational rejection, and facial expression [26]. Once a philosophical opposition to meat has formed, a dislike for the sensory and inherent qualities of meat occurred in vegetarians as a way of supporting and internalizing meat avoidance [14]. These components of disgust were thought to motivate further avoidance of meat. Thus, it is expected that a complete hedonic shift from liking to disliking or being disgusted at meat is required to

maintain strict vegetarian practices. The most frequent reasons given for red meat abstinence were also dislike of the taste and the high-fat content of red meat [27]. The response of current vegetarians to the flavor of meat differed from those of former vegetarians and non-vegetarians [16]. These results suggest that a taste factor can act on meat consumption as a cause or a consequence.

Although food choices are influenced by a broad range of economic, social and behavioral variables (e.g., food availability), taste and other sensory properties of foods have been deemed a major determinant of the selection of one food over another [28]. Thus, it was expected in the present study that a sensory hedonic shift regarding meat among vegetarians would influence other food choices and ultimately affect weight status. The motivations for and the nutritional outcomes of meat avoidance have been studied extensively. However, little is known about how meat consumption behavior affects total food intake and body weight by interacting with taste perceptions.

Therefore, the present study investigated the interplay among meat consumption behavior, energy intake, and weight status in female long term vegetarians and non-vegetarians, with a focus on the role of taste perception. Sensitivity to the bitter compound 6-n-propylthiouracil (PROP) and self-reported food liking/disliking were assessed as measures of taste perception for the following reasons. First, the ability to taste a low concentration of PROP has been linked to a newly discovered taste receptor gene, TAS2R38, thus providing a phenotypic