Continuing Medical Education for Community and Hospital Physicians in Israel

DY`Y[` $F^{1,2^*}\tilde{z}$?Ud`Ub`8 $^3\tilde{z}$ HUbXYhYf`< $^{1,2^*}$ UbX'G\jUfhn a Ub'D^{1,2}

¹Department of Family Medicine, Siaal Research Center for Family Medicine and Primary Care, Ben-Gurion University of the Negev, Beer-Sheva, Israel

²Clalit Health Service Organisation, Health Care Services, Israel

³Department of Otolaryngology, Head and Neck Surgery, Soroka University Medical Center, Beer-Sheva, Israel

*7cffYgdcbX]b[`Uih\cf. Prof. Roni Peleg, The Department of Family Medicine, Faculty of Health Sciences, Ben-Gurion University of the Negev, POB 653, Beer-Sheva, 84105, Israel, Tel: 972-8-6477436; Fax: 972-8-6477636; E-mail: pelegr@bgu.ac.il

FYWY] jY'XUhY. Jan 28, 2018; 5WWYdhYX'XUhY. Feb 08, 2018; DiV`]g\YX'XUhY. Feb 12, 2018

7 cdmf] [\h. © 2018 Peleg R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

5VghfUWh

6UW_[fc i bX. In this study we aimed to evaluate Continuing Medical Education (CME) activities of primary care physicians.

AYh\cXg. An e-mail questionnaire, based on the Jefferson Scale of Physician Life Long Learning (JSPLL), was sent to hospital- and community-based physicians in the Negev District.

FYg i hg. Eighty eight hospital-based physicians (38% response rate) and 118 community-based physicians (31% response) participated in the study. Hospital physicians participated more in annual meetings of professional medical associations (P<0.0001). They also gave more talks at professional meetings and took part more in research studies (P<0.0001). The variables that predicted the upper tertile of the most updated physicians were age (OR=0.947; 95% CI 0.926-0.968), male gender (OR=2.502; 95% CI 1.019-4.268), and active participation in teaching (OR=4.013; 95% CI 1.347-11.954).

7 cbW ig]cb. Both hospital and community physicians have a high level of access to medical journals and the Internet, but CME activities have been adapted more by hospital physicians.

?YmkcfXn '7~hHbi]b[a YX]W YXi Whcb/7ca a i b]hmLbX\cgd]HU VLgYX* ````cb[a YX]W YXi Whcb

Ir

Study population

 $\label{eq:started_st$

Comparison between the study groups in terms of reading habits and literature updates

Attitudes towards CME

Y>Y Yrson Scale of Physician Lifelong Learning (JSPLL)

a Yub'gWafY cb'h Y
 D@@ei YgJcbbUJfY k Ug'
 [[\Yf'Ua cb[\cgd]]U'd\ng Wubg'h Ub'Wa a i b]
hnid\ng Wubg'ff("- '†', "&
jg'')-"+'† +'%
dOS'SSS%L" ``]Z'cb[``YUfb]b[`fi b[``YUf

	No	26 (40.0)	2 (3.6)	
	Yes	13 (21.3)	46 (83.6)	<0.0001
Did you participate in the development of teaching programs?	No	48 (78.7)	9 (16.4)	
For PCPs			;	
	Urban	27 (61.4)	13 (72.2)	0.417
Location of main place of work	Other	17 (38.6)	5 (27.8)	
	Clinic director	6 (13.6)	11 (61.1)	<0.0001
Position in CHS	Primary physician/other	38 (86.4)	7 (38.9)	
	Family Medicine	19 (42.2)	6 (33.3)	0.514
Specialty	Other	26 (57.8)	12 (66.7)	
	Mean ± SD	43.4 ± 17.8	32.4 ± 13.8	0.027
Average number of patients in a work day (7 hour)	Range	9-100	Jul-50	

Upper tertile: LL activity score of 67-76. Bottom tertile: LL activity score of 41-58; JSPLL: Jefferson Scale of Physician Life Long Learning; PCP: Primary care physician; CHS: Clalit Health Services