

Detecting Abnormalities and their Consequences for Chronic Renal Failure

Emmanuel Marques*

Department of Pathology, University of Pittsburgh, Pittsburgh, USA

***Corresponding author:** Emmanuel Marques, Department of Pathology, University of Pittsburgh, Pittsburgh, USA, E-mail: emmarq@UoP.edu

Received: 26-Feb-2024, Manuscript No. JCEP-24-130927; **Editor assigned:** 29-Feb-2024, PreQC No. JCEP-24-130927 (PQ); **Reviewed:** 14-Mar-2024, QC No. JCEP-24-130927; **Revised:** 21-Mar-2024, Manuscript No. JCEP-24-130927 (R); **Published:** 28-Mar-2024, DOI: 10.4172/2161-0681.24.14.491

Citation: Marques E (2024) Detecting Abnormalities and their Consequences for Chronic Renal Failure. J Clin Exp Pathol. 14:491.

Copyright: © 2024 Marques E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Description

An insufficient ability of the kidneys to filter waste materials from the blood is the sign of renal failure, sometimes referred to as kidney failure. Failure like this might put the body at risk for major health problems like toxin accumulation and electrolyte imbalances. Renal failure can be acute or chronic, with each form having distinct causes, symptoms, and treatment approaches.

Acute Renal Failure (ARF)

ARF occurs suddenly, often within hours to days, and is usually reversible if diagnosed and treated promptly. Common causes include