Research Article Open Acces

*Corresponding author: Sharad Khandelwal R, Department of Microbiology, Institute of Life Sciences, HPT Arts and RYK Science College, Afýliated to Savitribai Phule Pune University, Nashik, India, Tel: 09689410568; E-mail: baviskarjayashree@gmail.com

Received February 11, 2016; Accepted March 14, 2016; Published March 21, 2016

Citation: Jayashree T 2 ā 16 l ƙ , CienAr ᴅ ʊ l c l Ж

was kept for 15 min. and made up to the mark with distilled water. Absorbance was measured at 553 nm against a reagent blank, prepared in same way as above [6,7].

Isolation and Identification of Arsenic oxidizing microorganisms from effluent: Out of collected industrial effluents, 04 different types of samples were chosen for bioremediation studies such as dye effluent, Coal Ash, Pesticide sample and chrome plating effluent. Enrichment of culture was done by adding 1% of e uent to nutrient broth containing Arsenite 50, 100, 150, 200, and 250 ppm, and was incubated at 28°C (Table 1). Microbial Culture from enriched asks was plated on nutrient agar plates with increasing Arsenic concentrations. Isolates with con uent growth were identi ed from Bac-Test Laboratory, Nasik [8].

Biomass determination

Isolates along with Pseudomonas putida in the e uents and in the MSM medium containing standard arsenic of 250 ppm were grown [8,9]. e growth of the microorganisms was determined a er every 24 hrs by taking absorbance at 545 nm.

Bioremediation studies

Bioremediation of Arsenic was done using di erent isolates, standard strain of Pseudomonas putida, in Erlenmeyer ask containing sterile 50 ml double strength MSM media (Nacl=5 gm/L, mgCl =0.5 gm/L, KH PO =0.45 gm/L, K HPO =0.9 gm/L, NH Cl=0.3 gm/L, KCl=0.3 gm/L, Glucose=1 gm/L, Yeast extract=1 gm/L) and sterile 50 ml e uents. Beside this 250 ppm/ ml of standard Arsenic (III) was also considered for bioremediation studies using all the isolates [10]. Oxidation was measured a er every 48 hrs spectrophotometrically [7]. Bioremediation of Arsenic was also performed using consortium Klebsiella pneumonia+P. putida and Lysinibacillus fusiformis+Bacillus subtilis.

Biosorption studies

Bacterial cells from actively growing culture were harvested and dry mass of 0.1 gm/10 ml of MSM was used for biosorption studies. Biomass was suspended in a ask containing Arsenic concentration of 250 ppm/ml. e binding capacity of the cells was calculated from the initial and nal concentrations [11].

Arsenic removal by Biosorption and Phytoremediation

1% of water hyacinth extract were added to MSM medium containing 250 ppm/ml of Arsenic, 1% of biomass was inoculated and Arsenic concentration was determined spectrophotometrically a er every 24 hrs [11].

Biosorption kinetics studies

e metal biosorption (q) and bioremoval e ciency (R) using bacterial biomass and plant extract was calculated using following formulae.

$$q = \frac{(C_i - C_t)}{M} V$$

$$R(\%) = \frac{(C_i - C_f)}{C_i} 100$$

Where: q=Metal adsorption (mg/g); M=dry biomass; V=total volume of sample treated; R=bioremoval e ciency %; C_i=initial concentration of metal in aqueous solution (ppm/L) and C_f=final concentration of metal in aqueous solution (ppm/L).

Toxicity testing on plants

Toxicity of Arsenic (III) and As (V) was observed on fenugreek plants with various dilutions of standard arsenic and di erent parameters root length, shoot length, size of leaves of fenugreek plant were determined.

Detoxi cation studies of e uents

Culture of Bacillus subtilis was inoculated in sterile Nutrient broth and incubated at room temperature for 24 hrs. to get cell density upto 10^6 cells/ ml. Detoxi cation of treated and untreated e uent against culture of Bacillus subtilis was studied on Mueller Hinton medium using Kirby Bauer's well di usion method. Distilled water was considered as control [12].

Results and Discussion

Sample analysis

Determination of pH and di erent parameters of industrial e uents, pH of all e uents was lower than neutrality i.e., 6.0, Pesticide e uent was found to be neutral around 7.1 (Figure 1).

Isolation and identi cation of Arsenic oxidizing organisms

Arsenic oxidizing bacteria have been identi ed from various arsenic rich environments [13]. Microorganisms were isolated on plates containing increasing Arsenic concentrations of 50, 100, 150, 200, and 250 ppm and the results as per Bac-Test Laboratory (Nasik, India) were as follows:

We could isolate four e ective microorganisms from various sources viz. Dye e uent, Chrome plating e uent, Coal ash sample, and pesticide contaminated soil, at a maximum range of arsenic concentration of 250 ppm tolerable by these organisms.

e salts were sterilized and inoculated with the respective isolates, (Isolate I- Dye isolate, Isolate II- Chrome isolate: Isolate III- Coal ash isolate: Isolate IV- Pesticide isolate) For pure 250 ppm of Arsenic, highest removal i.e., 81% was obtained by Isolate IV followed by isolate I i.e., 80%, Isolate III of 77%, followed by Isolate II i.e., 30% and Pseudomonas putida 76%. e samples were analyzed at the intervals of two days upto 10 days respectively (Graph 3a).

Arsenic, removal from e uents was obtained by Isolate III i.e., 75% followed by isolate I (37.35%) for concentration of 249 ppm and isolate II (60%) for concentration of 120 ppm in 10 days at room temperature (Graph 3b).

Bioremediation of e uent was also carried out using Standard strain of Pseudomonas putida which showed 83.1%, 75.3% and 48% removal of Arsenic from Coal ash e uent, Dye e uent and Chrome Plating e uent respectively (Graph 4).

Bioremediation of arsenic by consortium

Using the concentration of 250 ppm, Arsenic removal was obtained upto 92.8% by consortium of Klebsiella pneumoniaP and putida and around 94% by consortium of Lysinibacillus fusiformis and Bacillus subtilis (Graph 5).

Biosorption

Biosorption of As is studied using fungal biomass, and also fresh water micro algae [16]. In our study dry bacterial biomass of the isolates was used. Where 1% dry biomass showed signi cant Arsenic biosorption of 58 mg/g, 151 mg/g and 22 mg/g by isolate III, isolate II and isolate I with 8.8%, 60.4% and 23.2% bioremoval e ciency