

Early Perinatal Palliative Care Engagement can Assist to Improve Treatment During Pregnancy

James Baker*

Abstract

A rising variety of complicated foetal disorders that can be identified during pregnancy provide challenges for the newborn both before and after birth. It might be difficult for doctors who provide counseling to families in these situations to decide which ones would benefit from an early referral to palliative care. We argue that regardless of the offer chances for care planning that takes into account how to respect and honour the life of the foetus or baby.

*Corresponding author:

Received:

Editor assigned:

Revised:

Reviewed:

Published:

Citation:

Copyright:

Proportionality, terminal suffering and the
value of life. This paper explores the ethical
implications of the proportionality principle in
end-of-life care. It argues that the principle
requires a balance between the benefits and
burdens of treatment. In the context of
terminal illness, this balance is often
difficult to achieve. The paper discusses the
role of palliative care in alleviating suffering
and the importance of patient autonomy in
decision-making. It concludes that the
proportionality principle is a valuable tool
for guiding ethical decisions in end-of-life
care.

The proportionality principle is a key ethical
concept in end-of-life care. It requires that
the benefits of a treatment or intervention
be weighed against the burdens. In the
context of terminal illness, this often means
balancing the potential for prolonging life
with the risk of increased suffering. The
principle is particularly relevant in cases
where the patient's quality of life is severely
impaired. It provides a framework for
making difficult decisions about the best
course of action for the patient.

Proportionality is a central ethical principle
in healthcare. It is particularly important
in the context of end-of-life care, where
the goal is to provide the best possible
quality of life for the patient. This involves
weighing the benefits of treatment against
the burdens. The principle is often used to
guide decisions about the use of life-sustaining
treatment. It is a complex issue, and
there is no simple answer. However, the
proportionality principle provides a useful
framework for thinking about these issues.

Proportionality, terminal suffering and the
value of life. This paper explores the ethical
implications of the proportionality principle in
end-of-life care. It argues that the principle
requires a balance between the benefits and
burdens of treatment. In the context of
terminal illness, this balance is often
difficult to achieve. The paper discusses the
role of palliative care in alleviating suffering
and the importance of patient autonomy in
decision-making. It concludes that the
proportionality principle is a valuable tool
for guiding ethical decisions in end-of-life
care.

C
The proportionality principle is a key ethical
concept in end-of-life care. It requires that
the benefits of a treatment or intervention
be weighed against the burdens. In the
context of terminal illness, this often means
balancing the potential for prolonging life
with the risk of increased suffering. The
principle is particularly relevant in cases
where the patient's quality of life is severely
impaired. It provides a framework for
making difficult decisions about the best
course of action for the patient.

A
C
The proportionality principle is a key ethical
concept in end-of-life care. It requires that
the benefits of a treatment or intervention
be weighed against the burdens. In the
context of terminal illness, this often means
balancing the potential for prolonging life
with the risk of increased suffering. The
principle is particularly relevant in cases
where the patient's quality of life is severely
impaired. It provides a framework for
making difficult decisions about the best
course of action for the patient.

References

Proportionality, terminal suffering and the
value of life

Wool C, Cote-Arsenault D, Perry Black B, Denny-Koelsch E, Kim S, et al.
Wool C (2013)
Wool C, Parravicini E (2020)
Wilkinson D, de Crespigny L, Xafs V (2014)

Citation:

Viallard ML, Moriette G (2017)

Kersting A, Wagner B (2012)

Creating evidence: findings from a

Wool C, Catlin A (2019) Perinatal bereavement and palliative care offered
