

## Abstract

Potato is threatened by several soil-borne fungi causing wilt and root rots. In this study, two fodder radish (*Raphanus sativus* L.) (FR) cultivars (cvs. Boss and Defender), used as green manure preceding a potato crop, were evaluated for their suppressive effects against wilt incidence and severity, potato growth and yield as compared to animal manure. The essay was carried out in a completely randomised design with three types of organic amendment and two potato cultivars (cvs. Spunta and Royal). Incidence of potato wilting noted 100 days post planting (DPP) was high, exceeding 70%, for all soil amendments tested. The extent of vascular discoloration varied depending on amendments used wAreen manure preceding a potati1(fecm2rng )-1Span kaLang potybehavto crowilti, which6(control7 BDC, which6(ment

---

**\*Corresponding author:** H. Jabnoun-Khiareddine, UR13AGR09- Integrated Horticultural Production in the Tunisian Centre- East, Regional Center of Research on Horticulture and Organic Agriculture, University of Sousse, 4042, Chott-Mariem, Tunisia, Tel: 0021673327543; E-mail: [jkhayfa@yahoo.fr](mailto:jkhayfa@yahoo.fr)


**Received** February 02, 2016; **Accepted** February 13, 2016; **Published** February 19, 2016

**Citation:** Jabnoun-Khiareddine H, Aydi Ben Abdallah R, Ayed F, Gueddes-Chahed M, Hajlaoui A, et al. (2016) Effect of Fodder Radish (*Raphanus sativus* L.) Green Manure on Potato Wilt, Growth and Yield Parameters. *Adv Crop Sci Tech* 4: 211. doi:[10.4172/2329-8863.1000211](https://doi.org/10.4172/2329-8863.1000211)

**Copyright:** © 2016 Jabnoun-Khiareddine H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Phosphorus, Fertilizer, Vegetable, Rhizobia, Sclerotinia

---





Citation:

---



**Citation:** Jabnoun-Khiareddine H, Aydi Ben Abdallah R, Ayed F, Gueddes-Chahed M, Hajlaoui A, et al. (2016) Effect of Fodder Radish (

---

**Citation:** Jabnoun-Khiareddine H, Aydi Ben Abdallah R, Ayed F, Gueddes-Chahed M, Hajlaoui A, et al. (2016) Effect of Fodder Radish (*Raphanus sativus* L.) Green Manure on Potato Wilt, Growth and Yield Parameters. *Adv Crop Sci Tech* 4: 211. doi:[10.4172/2329-8863.1000211](https://doi.org/10.4172/2329-8863.1000211)

---

56. Molina OI, Tenuta M, El Hadrami A, Buckley K, Cavers C et al. (2014) Potato early dying and yield responses to compost, green manures, seed meal and chemical treatments. *Am J Potato Res* 91: 414-428.
  57. Morra MJ, Kirkegaard JA (2002) Isothiocyanate release from soil-incorporated Brassica tissues. *Soil Biol Biochem* 34: 1683-1690.
  58. Kirkegaard JA, Sarwar M (1998) Biofumigation potential of brassicas I. Variation
  59. Kirkegaard JA, Wong PTW, Desmarchelier JM (1996) In vitro suppression of fungal root pathogens of cereals by Brassica tissues. *Plant Pathol* 45: 593-603.
-