Efficacy of Ceftolozane/Tazobactam in Infections with Multidrug-Resistant Gram-Negative Bacteria: A Preliminary Clinical Experience in a Highly Endemic Hospital

Lucia Brescini[1,2*], Sara Mazzanti[1,2*], Luca Battistelli[2], Gianluca Morroni[2], Francesco Pallotta[1,2], Bianca Candelaresi[1,2], Silvia Olivieri[1,2], Francesco Ginevri[1,2], Giulia Cesaretti[1,2], Raffaella Donati[1], Matteo Cimatti[1], Rosaria G Polo[3], Vanessa Di Muzio[3], Michele Di Prinzio[3], Oriana Cornacchini[4], Marcello Tavio[1], Andrea Giacometti[1,2], Abele Donati[2,5], Elisabetta Cerutti[4] and Francesco Barchiesi[1]

[1]*Infectious Diseases Clinic, Azienda Ospedaliero University Delle Marche, Ancona, Italy*
[2]*Department of Biomedical Sciences and Public Health, Polytechnic University of Marche Medical School, Ancona, Italy*
[3]*Hospital Pharmacy, Azienda Ospedaliero University Delle Marche, Ancona, Italy*
[4]*Anesthesia, Transplant and Surgical Intensive Care Unit, Azienda Ospedaliero University delle Marche, Ancona, Italy*
[5]*Clinic of Anesthesia and General Resuscitation, Respiratory and Major Trauma, Azienda Ospedaliero University delle Marche Ancona, Italy*

Abstract

Introduction: Ceftolozane/Tazobactam is a new cephalosporin/beta-lactams inhibitor combination proven to be a drug with efficacy against urinary tract infections, abdominal infections and nosocomial ventilator-associated pneumonia, supported by multi-resistant Gram-negative

Keywords: Multi drug-resistant bacteria; Ceftolozano-tazobactam; *Pseudomonas*; ICU

Abbreviations: ICU: Intensive Care Units; IQR: Interquartile Range; MALDI-TOF/MS: Matrix-Assisted Laser Desorption/Ionization Time of Flight Mass Spectrometry; CCI: Charlson Comorbidity Index; HAP:

***Corresponding author:** Dr. Lucia Brescini, Infectious Diseases Clinic, Azienda Ospedaliero University Delle Marche, Ancona, Italy, E-mail: l.brescini@staf.univpm.it

Dr. Sara Mazzanti, Infectious Diseases Clinic, Azienda Ospedaliero University delle Marche, Ancona, Italy, E-mail: sara.mazzanti@hotmail.it

Received: 25-Jul-2023, Manuscript No. JIDT-23-108056; **Editor assigned:** 27-Jul-2023, PreQC No. JIDT-23-108056(PQ); **Reviewed:** 11-Aug-2023, QC No. JIDT-23-108056; **Revised:** 18-Aug-2023, Manuscript No. JIDT-23-108056 (R); **Published:** 25-Aug-2023, DOI: 10.4172/2332-0877.1000558

Citation: Brescini L, Mazzanti S, Battistelli L, Morroni G, Pallotta F, et al. (2023) Efficacy of Ceftolozane/ Tazobactam in Infections with Multidrug-Resistant Gram-Negative Bacteria: A Preliminary Clinical Experience in a Highly Endemic Hospital. J Infect Dis Ther 11: 558.

Copyright: © 2023 Brescini L, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

(p 0.020) and ICU stay (p 0.009). To investigate the reason for the higher mortality of patients admitted to the ICU, a second analysis was conducted comparing patients in intensive care and non-intensive care. (Table 2). ere were no di erences in gender and age between two groups. ICU patients were characterized by multiple serious comorbidities. Cardiovascular diseases, chronic renal failure, neutropenia and GI tract diseases were signi cantly more present in the group of patients in ICU (with p 0.046, p 0.031, p 0.041 and p 0.019, respectively). Even acute comorbidities with a strong impact on mortality, such as septic shock, were statistically more present in patients admitted to the ICU (p 0.031). As regards factors predisposing infection, patients in intensive care were more o en carriers of devices such as CVC ($p<0.001$), SNG ($P<0.001$), CV ($p<0.001$), surgical drainage ($p<0.001$). Moreover, as can be imagined, patients in the ICU were subjected to mechanical ventilation with a statistically signi cant di erence compared to the others ($p<0.001$). In addition, CVVH was more frequently used in ICU (p 0.01) and patients in intensive care department were more frequently post-surgery patients (p 0.004). Finally, a comparison of the two groups' shows that in ICU bacteria were isolated more frequently from BAL (p 0.046) and Ce olozano/ tazobactam was used less frequently in monotherapy (p 0.005). Signi cantly variables con rmed ad multivariate analysis were the presence of CVC (p 0.021), SNG (p 0.030), concomitant infections (p 0.043) and monotherapy (p 0.047).

Citation: Brescini L, Mazzanti S, Battistelli L, Morroni G, Pallotta F, et al. (2023) E cacy of Ceftolozane/ Tazobactam in Infections with Multidrug-Resistant Gram-Negative Bacteria: A Preliminary Clinical Experience in a Highly Endemic Hospital. J Infect Dis Ther 11: 558.