Case Report Open Access

**Ana Isabel Cobo-Cuenca* and Noelia Martín-Espinosa**

*Department of Nursing and Physiotherapy, University of Castilla La Mancha (UCLM), Spain*

Keywords:End stage care; Palliative care; Congestive heart failure

## Introduction

An aging population and a high rate of coronary heart disease and high blood pressure have led to an increase in the prevalence of chronic heart failure in Spain [1].

Deaths resulting from heart failure are as common, if not more common, as deaths resulting from certain types of cancer, and they o en occur as a result of the development of cardiac disease [2].

e illness is considered terminal when advanced heart failure is refractory to medical therapy and a heart transplant cannot be administered. Patients su er from severe shortness of breath and the persistence of symptoms such as congestive heart failure, and remain unresponsive to diuretic drugs, vasodilators, etc. [3,4].

According to the World Health Organization (WHO), palliative care is a new approach aimed at improving the quality of the lives of patients and their families when confronted with the complications associated with a life-threatening illness. It aims to prevent and alleviate su ering by providing pain management as well as physical, psychosocial and spiritual support to the patient [5].

Palliative care does not hasten or delay death, but it provides a genuine system of support for the patient and their family.

e end-stageis the phasethat precedes death. Patients gradually arrive at this stage a erintense physical deterioration, extreme debilitation, cognitive disorders and delirium, di culty of nutrient intake, and a prognosis of days or hours of remaining life.

Patients usually enter the end stage when they are admitted to the emergency room (ER). For this reason, it is vital that ER medical personnel (doctors, nurses, psychologists) be suitably trained to treat patients and their families [6].

e objective of this article is to present a case that introduces a palliative care plan for an 87-year-old male end stage patient arriving at the emergency room with acute pulmonary edema and congestive heart failure by ischemic cardiomyopathy.

***Corresponding author:** Cobo-Cuenca AI, Department of Nursing and Physiotherapy, University of Castilla la Mancha (UCLM), Toledo, Spain, E-mail: anaisabel.cobo@uclm.es

**Received** July 24, 2013; **Accepted** August 28, 2013; **Published** August 31, 2013

**Citation:** Cobo-Cuenca AI, Martín-Espinosa N (2013) End of Life in Stage Heart Failure in Elderly Individuals. J Palliative Care Med 3: 157. doi:10.4172/2165-7386.1000157

**Copyright:** © 2013 Cobo-Cuenca AI, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.