Abstract

Root-knot nematode disease caused by *Meloidogyne incognita* (Kofoid and White) Chit wood**)** is one of the major constraints for successful cultivation of tomato (*Lycopersicon esculentum* Mill.) in Ethiopia. Hence, the present study was conducted to evaluate the effect of leaf and seed extracts of four botanicals viz., Rape seed (*Brassica napus* L.)*,* Lantana (*Lantana camara* L.*),*

juveniles of M. incognita in vitro at the 10% concentration after 72 h of exposure time. There were no significant differences among the treatments of rape seed leaf (84.7%) at 10% concentration and

***Corresponding author:** Thangavel Selvaraj, Department of Plant Sciences, College of Agriculture and Veterinary Sciences, Ambo University, Ambo, Post Box No: 19, Ethiopia, East Africa Tel:+251-913073294; E-mail: tselvaraj_1956@yahoo.com

Received June 09, 2015; **Accepted** November 14, 2015; **Published** November 20, 2015

Citation: Feyisa B, Lencho A, Selvaraj T, Getaneh G (2015) Evaluation of Some Botanicals and *Trichoderma Harzianum* for the Management of Tomato Root-knot Nematode (*Meloidogyne Incognita* (Kofoid and White) Chit Wood). Adv Crop Sci Tech 4: 201. doi:10.4172/2329-8863.1000201

Copyright: © 2015 Feyisa B, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

konaensis. J Nematol 26: 744-748.

49. Gomez KA, Gomez AA (1984) Statistical Procedures for Agricultural Research, 2nd Edn, John Willey and Sons, New York, 680.

50. SAS Institute (2002) The SAS system for windows, version 9.1. SAS, Institute, Cary, NC.

51. Agbenin NO, Emechebe AM, Marley PS (2004) Evaluations of neem seed powder for Fusarium wilt and Meloidogyne control on tomato. Arch. Phytopathol. Plant Protection 37: 319-326.

52. Parmar BS (1987) An overview of neem research and use in India during the years 1983-1986; in: Natural Pesticides from the Neem Tree and other Tropical Plants; Proceedings of the 3rd International Neem Conference, Nairobi (1996), (eds) Schmutterer H, Ascher KRS.;; GTZ Eschborn, Germany 55–80.

53. Elbadri GAAD, Lee W, Park JC, Yu HB, Choo HY (2008) Evaluation of various