

## Evidence Based use of Heat, Cold and NSAIDS for Plantar Fasciitis

Jerrold S. Petrofsky<sup>1\*</sup>, Michael S. Laymon<sup>2</sup>, Faris Alshammari<sup>1</sup> and Iman Akef Khowailed<sup>1</sup>

<sup>1</sup>Department of Physical Therapy, Loma Linda University, Loma Linda, California, USA

<sup>2</sup>School of Physical Therapy, Touro University Nevada, Henderson, USA

\*Corresponding author: Dr. Jerrold Petrofsky, Professor and Director of Research, Department of Physical Therapy, Loma Linda University, Loma Linda, California 92350, USA, Tel: 909558 7274; E-mail: [jpetrofsky@llu.edu](mailto:jpetrofsky@llu.edu)

Received date: Mar 10, 2014, Accepted date: May 26, 2014, Published date: June 05, 2014

Copyright: © 2014 Petrofsky JS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

Plantar Fasciitis is a possible complication in runners and even people who walk a lot. There have been numerous proposed therapies to treat this disorder but the independent effect of heat, cold and NSAIDS have not been examined without the concomitant use of other modalities. Here we examined the effect of heat, cold and Advil independently of other therapies on plantar fasciitis.

**Objective:** Subjects with plantar fasciitis either had no intervention, cold applied 20 minutes at night before bed, or 20 minutes in the morning upon waking or either moist heat at night or in the morning (1 hour) or dry heat (4 hours) at night for 1 day or 3 days with and without Advil. Plantar fascia swelling, tenderness and pain were evaluated.

**Methods:** Visual analog pain scale, plantar facial thickness was measured by ultrasound as a measure of inflammation, the pressure tolerated by force on the plantar fascia from an algometer were measured first thing in the morning. An activity of Daily Living Subscale of the Foot and Ankle Ability Measure (FAAM-ADL) questionnaire was used for assessing disability. There were 99 subjects in 9 groups. All post intervention measurements were taken first thing in the morning before activity.

**Results:** The greatest relief of symptoms was with the application of cold used at bedtime the night before the measurements, cold used in the morning was not as effective as was heat. Cold use reduced the thickness of the plantar fascia and pain. Cold plus Ibuprofen were significantly better at reducing plantar fascia symptoms than cold alone. Conclusion: Cold applied for 20 minutes prior bedtime was the most effective treatment for reduced symptomology caused by plantar fascia inflammation. Advil further reduces pain when used with cold.

**Keywords:** Cold temperature; Plantar fasciitis; Pain; Foot; Therapeutics

### Introduction

Plantar fasciitis is a common condition and can occur as a single occurrence or repeatedly throughout life [1,2]. It is most common in people who exercise frequently or especially in people with a high BMI or in the elderly [1,3,4]. The plantar fascia is a thick fibrous connective tissue which originates at the medial tuberosity of the calcaneus and inserts into the proximal phalanges [5,6]. The central portion is the thickest, attaches at the posterior aspect of the medial tuberosity of the calcaneus posterior to the origin of the flexor digitorum brevis tendon, and is 1.5 to 2.0 cm in width, distally, at the level of the metatarsophalangeal joints. The central portion of the plantar aponeurosis divides into five fascicles, one for each of the toes [5,7]. The lateral portion of the plantar aponeurosis arises from the lateral aspect of the medial tuberosity of the calcaneus and its distal medial and lateral bands attach to the plantar plate of the fourth toe and to the base of the fifth metatarsal respectively and is 1.0 to 1.5 cm in width. The medial portion of the plantar aponeurosis is thin and continuous medially with the dorsal fascia and laterally with the central portion of the plantar aponeurosis. Edema in the plantar fascia therefore affects movement in the foot by inhibiting normal tendon movement to flex

and extend the foot especially when body weight is applied [6,7]. Many mixed modalities have been used to treat this disorder. These include custom orthotic devices [7], rest, ice baths [5,8], heat, and stretching [7]. Night splinting and corticosteroid injections have also been used [9]. Heat has been applied at night and in many cases has positive results but the results are unpredictable [10]. But heat and other modalities are never used alone and also include other therapeutic modalities. But even with these modalities, the inflammation can return and severely limit normal gait [11].

One problem with treatment is the lack of evidence basis in the modalities that are used for treatment. Treatment involves mixed modalities [11,12]. Usually a physician or physical therapist will manipulate the foot by stretching and apply heat or cold and use non-steroidal anti-inflammatories or steroids and orthotics [12-14]. In some cases, platelet rich plasma has been injected [6]. If inflammation and pain are resolved in a clinical setting there is no way of knowing which modality resulted in improvement.

In the present investigation we used cold, heat or Advil as the only intervention for up to 3 days and measured plantar fascia inflammation, pain level, tolerance to pressure on the plantar surface, and a standardized functional assessment instrument. No other treatments such as stretching or range of motion or steroids were used the week before the subjects participated in this project. Two series of

experiments were undertaken. In the first, either heat or cold was used in an acute 1 day study where heat or cold were applied either the night before or morning before measurements. This led to the second series of experiments where cold and Advil were tested over a 3 day period to test their joint efficacy.

## Subjects

Ninety nine subjects were recruited for this study in 2 series of experiments. In one group, cold and heat were used for a single treatment and the other cold was used in conjunction with an NSAID,

Advil. Eligible subjects were required to meet all the following inclusion criteria: (1) subjects of both genders are diagnosed with plantar fasciitis; (2) subject's age is between 18-65 years; (3) duration of symptoms is less than 5 years; and, (4) plantar fascia thickness greater than 4 mm as assessed by imaging ultrasound. Exclusion criteria included: (1) previous fracture or surgery to the foot or (2) specific metabolic and connective tissue disorder associated with or contributing to the diagnosis of plantar fasciitis (PF). All purposes and procedures were approved by the Solutions Institutional Review Board and each patient signed a statement of informed consent. The demographics of the subjects are listed in Tables 1 and 2.

		Age (years)	Height (cm)	Weight (kg)	BMI	Faam
<b>Control</b>	<b>Mean</b>	51.9	155.2	85.4	29.4	30.2
	<b>Standard deviation</b>	3.6	26.5	15.8	4.2	8.5
<b>Heat night</b>	<b>Mean</b>	45.6	172.2	80.0	26.9	26.0
	<b>Standard deviation</b>	11.9	8.7	17.5	5.4	14.7
<b>Moist night</b>	<b>Mean</b>	46.6	176.6	86.8	27.7	23.2
	<b>Standard deviation</b>	6.4	12.7	17.7	4.3	13.2
<b>Moist morning</b>	<b>Mean</b>	42.8	167.3	80.3	28.7	23.8
	<b>Standard deviation</b>	9.0	3.8	15.3	5.2	16.9
<b>Cold night</b>	<b>Mean</b>	52.3	164.1	75.8	27.9	21.9
	<b>Standard deviation</b>	5.5	4.5	19.1	5.8	5.4
<b>Cold morning</b>	<b>Mean</b>	43.8	171.4	81.3	27.5	23.8
	<b>Standard deviation</b>	9.5	6.7	19.0	5.2	16.9



the medial calcaneal tubercle. The algometer contact head was aligned perpendicularly to the skin and the principal investigator gradually increased the algometer pressure until the patient reported pain. The algometer reading which represented the pressure needed to elicit pain (in Newton per square centimeter), was recorded. Higher algometer scores indicated greater pressure tolerance and, hence, less tenderness. Lower algometer scores indicated less pressure tolerance and, thus, greater tenderness. The reliability and validity of pressure algometer as a measure of tenderness has been documented in the literature.



Figure 1: A typical subject before modalities.




Figure 2 Percent change in the analog visual pain scale from the first to the second day after application of each modality. The data is the mean of 10 subjects +/- the SD.





Figure 3 Change in plantar fascia thickness pre and post modality. The data is the mean of 10 subjects +/- the SD.




**Figure 4** the change in algometer pressure for perception of pain before and after each modality in 10 subjects +/- the SD.

**Series 2** - Based on the series 1 experiments, only cold was tested in series 2 since this was the best modality of those we tested. It was tested for 3 days with and without Advil to see if an anti-inflammatory plus cold would have a better effect.

### Analog Visual Pain Scale

The results of the analog visual pain scale on the 3 groups of subjects are shown in Figure 5



cold and 18.7% for the cold plus Ibuprofen groups. There was no significant difference between the groups. The sham group showed an increase averaged 4.1% and was not significant compared to the resting value but there was no difference between the groups ( $p > 0.05$ ).




Figure 7: The average plantar thickness over the 4 days in the 3 groups of subjects. Each point is the mean +/- the SD.




Figure 8: The average FAAM score over the 4 days in the 3 groups of subjects. Each point is the mean +/- the SD.

## Discussion

Plantar fasciitis is a common disorder especially in runners [17]. Obesity, excessive foot pronation, excessive running, and prolonged standing are also risk factors for developing plantar fasciitis even in non-runners and non-athletes [7]. It can be diagnosed from ultrasound imaging of the plantar fascia if it is thicker than 4 mm [7]. Many different treatment modalities are used for this disorder. The most common treatment is stretching of the gastrocnemius and soleus muscles to loosen the plantar fascia [1,17]. Deep myofascial massage and iontophoresis is usually performed by a physical therapist and can help [2]. Orthotics commonly are used to keep pressure off of the plantar fascia [14]. NSAIDs have been reported to have excellent reductions in pain and inflammation [18]. Cold is used for treatment in some cases but papers showing efficacy are rare. Ice on the plantar

fascia reduces sensation by blocking receptors [19]. It also reduces plantar swelling [20].

But in these studies, cold or heat were not used alone but in conjunction with other therapies, especially stretching. Here we looked at modalities alone. Clearly from the series 1 experiments, cold the night before measures of pain and inflammation was the clear winner. It reduced swelling and pain significantly. Cold in the morning worked but not nearly as well. Heat showed good results at night, especially moist heat. But heat applied at night in other studies has shown positive results but the results are sometimes unpredictable [10].

Therefore in the series 2 experiments, we used only cold at night but wanted to see if cold every night for 3 days would have a better effect. We also tested to see if Advil, an NSAID would work synergistically with cold. Here, for the cold only group, there was a 57.9% reduction in heel pain on the AV pain scale; a 170% reduction in the algometer pressure that could be tolerated, and plantar fascia thickness was reduced by 16.5%. Cold here was used for 3 repeated days. When cold and Ibuprofen were added, data was significantly better than the cold treatment and showed a considerable improvement in the pressure that could be tolerated by the algometer on the plantar fascia. To some extent this is not surprising since it shows that Ibuprofen is a good pain reducing analgesic and is synergistic with cold [21,22]. Ibuprofen works by moderate inhibition of COX-1 and COX-2 enzymes thereby inhibiting prostaglandin production in tissue. It therefore has benefits in reducing pain and inflammation [21,23]. Therefore it should have dual effects on plantar fasciitis; reducing swelling and reducing pain. The reduction in plantar fascia thickness was not greater with cold plus Ibuprofen; it would appear that the predominant effect on swelling here was cold while the effect of Ibuprofen alone was not studied. The visual analog pain scale did not improve better than that seen for cold alone but this is not surprising since they had no pressure on the plantar fascia when the pain scale was recorded vs. the algometer where pressure was applied to the plantar fascia. Here the 2 modalities together made a significant difference.

The best evidence for the efficacy of cold comes when comparing it to the sham cold and the sham Ibuprofen group. Here there was no change in plantar fascia thickness, a small reduction in pain but no change in pain related to applying pressure. Thus cold and cold plus Ibuprofen are good together to help plantar fasciitis. A problem with cold is that it has always been poorly defined in terms of what cold is and how long to apply it [24]. Sometimes it involves whole body immersion, contrast baths [25], or icing with the length of time modalities are applied quite variable [24,26]. Cooling causes a reflexive vasoconstriction of blood vessels, due to increased affinity of alpha-adrenergic receptors in cold blood vessels for norepinephrine in the vascular walls [27]. This should contribute to reducing inflammation. For the foot, where the plantar fascia is close to the surface, cold should be effective in penetrating in the 20 minute treatment here with-

## References