*Corresponding author: Godfred Nwosu Ezena, Ghana Ministry of Food and Agriculture, PO Box 43, Bawku, Bawku-Zebilla Road, Ghana, Tel: 011233266514910; E-mail: goddie191@yahoo.com (or) cakotsenmensah@ug.edu

Received June 29, 2016; Accepted July 20, 2016; Published July 21, 2016

Citation: Ezena GN, Akotsen-Mensah C, Fening KO (2016) Exploiting the Insecticidal Potential of the Invasive Siam Weed, Chromolaena odorata L. (Asteraceae) in the Management of the Major Pests of Cabbage and their Natural Enemies in Southern Ghana. Adv Crop Sci Tech 4: 230. doi:10.4172/2329-8863.

Lamiaceae, Euphorbiaceae, Combretaceae and Annonaceae. ey are common weed, shrub and tree species on and around farms (Devanand and Rani) and Amoabeng et al. [13,16]. Pests control using these local materials o er the farmer the opportunity to reduce production cost, as the plants o en grows in the wild and around farms and so can be obtained with little e ort and zero or minimal cost Amoabeng et al. [13].

However, some botanicals may adversely a ect bene cial insects Buss & Park- Brown, Dubey et al. [17,18]. Amoabeng et al. [13] con rmed this in a eld experiment using 30 g/L *C. odorata* to control DBM larvae among other botanicals. ey showed that although the 30 g/L *C. odorata* extract was e ective in reducing pest populations, this concentration was however detrimental to natural enemies and non-targeted organisms in the eld. Fening et al. [19] have also cautioned that higher concentrations of botanicals could possibly have some detrimental e ects on the natural enemy populations because their numbers reduce with increasing concentrations. Hence, the current research was conducted to optimize the concentrations of crude extract from Siam weed *C. odorata* for e ective control of the key pests of cabbage with minimal or no adverse e ect on the natural enemies of these pests and other non- target organisms.

Materials and Methods

Citation: Ezena GN, Akotsen-Mensah C, Fening KO (2016) Exploiting the Insecticidal Potential of the Invasive Siam Weed, *Chromolaena odorata* L. (Asteraceae) in the Management of the Major Pests of Cabbage and their Natural Enemies in Southern Ghana. Adv Crop Sci Tech 4: 230. doi:10.4172/2329-8863.1000230

Citation: Ezena GN, Akotsen-Mensah C, Fening KO (2016) Exploiting the Insecticidal Potential of the Invasive Siam Weed, *Chromolaena odorata* L. (Asteraceae) in the Management of the Major Pests of Cabbage and their Natural Enemies in Southern Ghana. Adv Crop Sci Tech 4: 230. doi:10.4172/2329-8863.1000230

Acknowledgements

We are grateful to the Technical staff, of the Entomology Unit of the Forestry and Horticultural Crops and Research Centre (FOHCREC), University of Ghana, Ghana.

References

1. Asare-Bediako E, Addo-Quaye AA, Mohammed A (2010) Control of Diamond back moth (Plutella xylostella) on cabbage (Brassica oleracea var capitata) using intercropping with Non-Host Crops. American Journal of Food Technology 5: 269-274.
2. Timbilla JA, Nyarko KO (2004) A survey of cabbage production and constraints in Ghana. Ghana Journal of Agricultural Science 37: 93-101.
3. Timbilla JA (1997) Integrated pest management of DBM. P xylostella (Lepidoptera: Plutellidae) in Ghana. Masters of Philosophy Thesis, KNUST - Kumasi, Ghana, p: 125.
4. Fening KO (2013) Improving Maternal and Child Health: The Role of Food Safety in the Developing World. In: Maternal-Child Health-Interdisciplinary Aspects within the Perspective of Global Health. Uwe Groß and Kerstin Wydra (eds.), Göttingen International Health Network (GIHN), Universitätsverlag Göttingen, Germany, pp: 53-61.
5. Norman C, Shealy MD (2007) Illustrated Encyclopedia of Healing Remedies. (online) PHD, Elements Book Inc. 160 North, Washington Street. Boston MA 02114. Available from: http://www.hcun.com/healthandhealing/kcabbage/indexhtm.
6. De Lannoy G (2001) Leafy Vegetables in Crop Production in Tropical Africa. Raemaekers RH (ed), Directorate General for International Co-operation, Brussels, Belgium, pp: 403-511.
7. 0RFKLDK 0% %DLGRR 3. 2ZXVX $N\DZ 0 ,QÀXHQFH RI GLIIHUHQW QXWULHQW applications on insects populations and damage to cabbage. Journal of Applied Bioscience 38: 2564- 2572.
8. Ninsin DK (1997) Insecticides use pattern and residue levels in cabbage cultivated within the Accra-Tema Metropolitan areas of Ghana. Master of Philosophy Thesis, Insect Science Programme. University of Ghana, Legon, p: 85.
9. Obeng-Ofori D, Ankrah DA (2002) Effectiveness of aqueous neem extracts for the control of insect pest of cabbage (Brassica oleracea var capitata L) in the Accra plains of Ghana. Agriculture and Food Science Journal Ghana