

Exploring the Various Surgical and Clinical Procedures in Pancreas Transplantation

Dorge Reyes*

Department of Transplantation Surgery, University of Washington Medical Center, USA

Introduction

A pancreas transplant is a surgical procedure in which a healthy pancreas from a deceased donor is placed into a person whose pancreas no longer functions properly. The pancreas is located behind the lower part of the stomach and plays a crucial role in producing insulin, a hormone that helps regulate the absorption of sugar (glucose) into cells. When the pancreas is unable to produce enough insulin, blood sugar levels can rise to unhealthy levels, leading to type 1 diabetes. Pancreas transplants are most commonly performed to treat type 1 diabetes, offering the potential for a cure. However, this procedure is typically reserved for those with severe complications from diabetes, as the risks and side effects of the transplant can be significant. In some cases, pancreas transplants may also be considered for treating type 2 diabetes or, morlow insulin resistance and low insulin production, pancreas transplant may be a treatment option. About 10 percent of all pancreas transplants are performed in people with type 2 diabetes [2].

Transplantation of a pancreas alone, including

Pancreas transplant alone

People with diabetes and early or no kidney disease may be candidates for a pancreas transplant alone (solitary pancreas transplant). A pancreas transplant surgery involves the placement of a healthy pancreas into a recipient whose pancreas is no longer functioning properly.

Combined kidney-pancreas transplant

Surgeons often may perform combined (simultaneous) kidney-pancreas transplants for people with diabetes who have or are at risk of kidney failure. Most pancreas transplants are done at the same time as a kidney transplant. The goal of this approach is to give you a healthy kidney and pancreas that are unlikely to contribute to diabetes-related kidney damage in the future [3].

Pancreas-after-kidney transplant

For those facing a long wait for both a donor kidney and a donor pancreas to become available, a kidney transplant may be recommended first if a living- or deceased-donor kidney becomes available. After you recover from kidney transplant surgery, you'll receive a pancreas

transplant once a donor pancreas becomes available.

Pancreatic islet cell transplant

During pancreatic islet cell transplantation, insulin-producing cells (islet cells) taken from a deceased donor's pancreas are injected into a vein that takes blood to your liver. More than one injection of transplanted islet cells may be needed. Islet cell transplantation is being studied for people with serious, progressive complications from type 1 diabetes. It may only be performed as part of a Food and Drug Administration-approved clinical trial. The pancreas is an organ about the size of a hand located in the abdomen in the vicinity of the stomach, intestines, and other organs. It lies behind the stomach and in front of the spine. The pancreas produces juices that help digest food and hormones such as insulin and glucagon that maintain optimal blood sugar levels and help the body to use and store energy from food [4]. A pancreas transplant is an organ transplant that involves implanting a healthy pancreas (one that produces insulin) into a person whose pancreas no longer can supply sufficient insulin to the body. The healthy pancreas comes from either a deceased donor or in the form of a partial pancreas from a living donor. A pancreas transplant offers a potential cure for type 1 or insulin-dependent diabetes. A successful pancreas transplant will eliminate the need for insulin injections, reduce or eliminate dietary and activity restrictions due to diabetes, and decrease or eliminate the risk of severe low blood sugar reactions. A pancreas transplant can also help manage the damage to other organs including the kidneys that may result from type 1 diabetes [5]. Pancreas transplants are primarily offered to persons with type 1 with severe kidney disease or other life-threatening consequences from uncontrolled glucose levels. Type 1 diabetes is caused by a loss or malfunction of the insulin producing cells, called pancreatic beta cells. Beta cells (β cells) are a type of cell found in the pancreatic islets of the pancreas. They make up 65-80% of the cells in the islets [6].

References

- Marcus U (2019) HIV infections and HIV testing during pregnancy, Germany, 1993 to 2016. *Euro surveillance* 24: 1900078.
- Bunn JY, Solomon SE, Miller C, Forehand R (2017) Measurement of stigma in

*Corresponding author: Dorge Reyes, Department of Transplantation Surgery, University of Washington Medical Center, USA, Tel: 158962314785, Email: reyedor456@gmail.com

Received: 01-Nov-2024, Manuscript No: jcet-25-160423; Editor assigned: 04-Nov-2024, PreQC No: jcet-25-160423 (PQ); Reviewed: 18-Nov-2024, QC No: jcet-25-160423; Revised: 22-Nov-2024, Manuscript No: jcet-25-160423 (R); Published: 30-Nov-2024, DOI: 10.4172/2475-7640.1000257

Citation: Dorge R (2024) Exploring the Various Surgical and Clinical Procedures in Pancreas Transplantation. *J Clin Exp Transplant* 9: 257.

Copyright: © 2024 Dorge R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

3. Al-Ani R, Al Obaidy A, Hassan F (2019) Multivariate analysis for evaluation the water quality of Tigris River within Baghdad City in Iraq. *Iraqi J Agric Sci* 50: 331-342.
 4. Blann KL, Anderson JL, Sands GR, Vondracek B (2009) Effects of agricultural drainage on aquatic ecosystems: a review. *Crit Rev Environ Sci Technol* 39: 909-1001.
 5. Boynton W, Kemp W, Keefe C (1982) A comparative analysis of nutrients and other factors influencing estuarine phytoplankton production. In *Estuarine comparisons* 69-90.
 6. Samet J, Dominici F, Currier I, Coursac I, Zeger S (2000) Fine particulate air pollution and mortality in 20 US cities, 1987-1994. *N Engl J Med* 343: 1742-17493.
 7. Soto CJ, John OP, Gosling SD, Potter J (2011) Age differences in personality traits from 10 to 65: Big Five domains and facets in a large cross-sectional sample. *J Pers Soc Psychol* 100: 330-348.
 8. Jang KL, Livesley WJ, Angleitner A, Reimann R, Vernon PA (2002) Genetic and environmental influences on the covariance of facets defining the domains of the five-factor model of personality. *Pers Individ Dif* 33: 83-101.
 9. DeYoung CG, Quilty LC, Peterson JB (2007) Between facets and domain: 10 aspects of the Big Five. *J Pers Soc Psychol* 93: 880-896.
 10. Gosling SD, Vazire S, Srivastava S, John OP (2004) Should we trust web-based studies? A comparative analysis of six preconceptions about internet questionnaires. *Am Psychol* 59: 93-104.
-