



Novel disease results in the release of cytokines that are

According to reports, Punjab, Pakistan's eedom, lost 6 billion

Vaccination, on the other hand, is the most e ective strategy to treat

In the expression of the protein, there are three multiple ways or

according to OIE norms . It was constant in control as they weren't taking antigen in the form of vaccine. As we've started giving vaccine to group 2 and 3 from day 0 that's why HI titer start adding in them.

These findings, still, should be interpreted with caution, and further exploration through contagion challenge assay is needed to increase our understanding of the mechanisms involved. Still, if the challenge infection trial is carried out, the same humoral vulnerable responses in groups 2 and 3 offer a hopeful suggestion of the defensive effectiveness.

References

1. Meng Y, Wang C, Zhang L (2019) Recent developments and highlights in allergic rhinitis. *Allergy* 74: 2320-2328.
2. Ihara F, Sakurai D, Yonekura S, Iinuma T, Yagi R, et al. (2018) Identification of specifically reduced Th2 cell subsets in allergic rhinitis patients after sublingual immunotherapy. *Allergy* 73: 1823-1832.
3. Wise SK, Lin SY, Toskala E (2018) International consensus statement on allergy and rhinology: allergic rhinitis-executive summary. *Int Forum Allergy Rhinol* 8: 85-107.
4. Klimek L, Mullol J, Hellings P, Gevaert P, Mösges R, et al. (2016) Recent pharmacological developments in the treatment of perennial and persistent allergic rhinitis. *Expert Opin Pharmacother* 17: 657-669.

5. Berger WE, Meltzer EO (2015) Intranasal spray medications for maintenance