Advances in Crop Science and Technology

Mekonnen et al., Adv Crop Sci Tech 2016, 4:6
DOI:10.4172/2329-8863.1000245

Research Article OMICS International

Abstract

To assess the integrated effect of planting pattern and low dose herbicide mixtures on weeds and growth, yield attributes and yields of cowpea, and to determine the economic feasibility of different weed management practices LQ FRZSHD D ¿HOG H[SHULPHQW ZDV FRQGXFWHG DW 6LULQND $JULFXOWXUDO and Sirinka in Northern Ethiopia during the 2014 main cropping season. There were 16 treatments comprising the combinations of two planting patterns (60 cm × 10 cm, 45 cm × 15 cm) and eight weed management practices (s-metolachlor 2.0 kg ha^{-1}, s-metolachlor at 1.0 kg ha^{-1}+hand weeding and hoeing 35 weeks after crop emergence (WAE), pendimethalin at 1.0 kg ha^{-1}+hand weeding and hoeing 5 WAE, s-metolachlor at 1.0 kg ha^{-1}+pendimethalin at 1.0 kg ha^{-1}, s-metolachlor at 1.0 kg ha^{-1}+pendimethalin at 0.75 kg ha^{-1}, s-metolachlor at 0.75 kg ha^{-1}+pendimethalin at 1.0 kg ha^{-1}, hand weeding and hoeing 3 WAE and weedy check. The treatments were laid out in factorial combination in a randomized complete block design with three replications. The highest number of pods per plant, number of seeds per pod, and hundred seed weight were obtained from the combination of s-metolachlor at 1.0 kg ha^{-1}+hand weeding 5 WAE along with 60 cm × 10 cm at Sirinka. Higher (3092 kg ha^{-1}) grain yield was recorded at Sirinka than at Jari (2714 kg ha^{-1}). The highest (53460 ETB ha^{-1} JURVV EHQH¿W ZDV REWDLQHG IURP hand PHWRODFKOR weeding and hoeing 5 WAE, followed by pendimethalin at 1.0 kg ha^{-1}+hand weeding and hoeing 5 WAE (46737 ETB ha^{-1}). Therefore, managing the weeds with the application of 1.0 kg ha^{-1} of s- metolachlor+hand weeding and hoeing 5 WAE along with 60 cm × 10 cm proved to be the most feasible practice. Alternate herbicides for the control of X. VWUXPDULXP LQIHVWHG ¿HOGV LQ WKH VWXG\ DUHD QHHGV WR EH H[SORUHG

Keywords: Broadleaved and grass weeds; Cowpea; Grain yield; Herbicide mixtures; Integrated management; Pendimethalin; s-metolachlor; Vigna unguiculata; Weed

Introduction

Cowpea (Vigna unguiculata (L.) Walp.) is one of the most important food grain legumes in the tropics, including Africa, which accounts for 64% of the world production [1]. West Africa represents the largest production zone with modest amounts emanating from the east African countries of Mozambique, Tanzania, Uganda and to some extent Ethiopia [1,2]. In addition to its importance in human food, cowpea is also useful for soil fertilization through symbiotic nitrogen xation and can be a major animal feed due to the quality of its leaves [3].

It is cultivated around the world primarily for seed, but also as a vegetable (for leafy greens, green pods, fresh shelled green peas, and shelled dried peas), as cover crop and for fodder [4]. In most African

nigrum. Similarly, Wilson [12] found that for every 100 kg dry weight of weeds, cowpea yield was reduced by about 208 kg ha^{-1}. In Ethiopia, one timely, early weeding at 25 days a er emergence resulted in 70% yield increase of common bean and up to 300% increase in cowpea compared to the no-weeding [13].

Di erent management practices should be employed to reduce yield loss due to weeds. Among those practices, integrated weed managemen (IWM) involves a combination of cultural, physical, chemical and

*Corresponding author: Getachew Mekonnen, College of Agriculture and Natural Resources, Mizan Tepi University, PO Box 260, Mizan Tepi, Ethiopia, Tel: +251912008038; E-mail: sibuhmekdes@gmail.com

Received November 05, 2016; Accepted November 14, 2016; Published November 18, 2016

Citation: Mekonnen G, Sharma JJ, Negatu LW, Tana T (2016) Growth and Yield Response of Cowpea (Vigna unguiculata L. Walp.) to Integrated Use of Planting Pattern and Herbicide Mixtures in Wollo, Northern Ethiopia. Adv Crop Sci Tech 4: 245. doi: 10.4172/2329-8863.1000245

Copyright: © 2016 Mekonnen G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

overall impact of weeds and, indeed, maximize the bene ts. e use of a single herbicide may result in shi of the weed ora in favour of the species that are not controlled, thus may increase the problem in the future.

Moreover, to manage mixed population of weeds and also to avoid herbicide resistance development by continuous use of a single herbicide, compatible mixtures can be employed to widen the spectrum of weed suppression. Herbicide combinations can give spectacularly good control at doses considerably below those normally applied in a single application. It may be additive or synergistic or prevent rapid detoxi cation of herbicides and are safer to crops than application of a single herbicide alone. e use of herbicide combinations is not new, but it has not received the attention and input that is necessary to fully understand and implement the practice. erefore, there is a need for evaluation of a range of herbicides alone and as a tank mixture to have broad spectrum weed management [15].

e present study, therefore, is intended 1) To assess the integrated e ect of planting pattern and low dose herbicide mixtures on weeds, nodulation, growth, yield attributes and yields of cowpea, and 2) to determine the economic feasibility of di erent weed management practices in cowpea.

Materials and Methods

Description of the study area

e experiment was conducted at Jari experimental sites (11°21'N latitude and 39°38'E longitude; 1680 masl. altitude) at Sirinka Agricultural Research Center and Sirinka (11°45'00" N latitude; 39°36'36"E longitude; 1850 masl altitude) in northern Ethiopia during the 2014 main cropping season. e soil of the experimental elds was clay loam and clay, while the pH was 6.98 and 6.94 at Sirinka and Jari, respectively. At Sirinka, the organic carbon was 1.35%, total N was 0.07%, available P 13.7 mg kg^{-1} soil and CEC 56.47 cmolC kg^{-1}, while the respective values at Jari were 1.37%, 0.05%, 11.17 mg kg^{-1} soil and 47.44 cmolC kg^{-1}. e total rainfall received during the crop season was 795.4 and 649.1 mm at Sirinka and Jari with mean maximum and minimum temperatures of 27.0 and 14.2°C, and 30.1°C and 16.0°C, respectively (Figure 1). Soil sample analysis was done at the Sirinka Agricultural Research Center.

Experimental materials

e cowpea variety Asrat (ITS 92KD-279-3), released by Sirinka Agricultural Research Center/Amhara Region Agricultural Research Institute (SRARC/ARARI) in 2001, was used in these experiments. e variety is well adapted to moisture stress areas in the northeast Wollo and similar lowland areas. is variety is suitable for an altitudinal range of 1450-1850 masl and annual rainfall of 660-1025 mm. It is bushy and trailing type I. It attains physiological maturity in 95-100 days [16]. Description of herbicides (s-metolachlor and pendimethalin) used in the experiment has been presented in tabular form hereunder Table 1.

Treatments and experimental design

ere were 16 treatment combinations comprising of two planting patterns (60 cm × 10 cm and 45 cm × 15 cm) and eight weed management practices (s-metolachlor at 2.0 kg ha^{-1}, s-metolachlor at 1.0 kg ha^{-1}+hand weeding and hoeing 5 weeks a er crop emergence (WAE), pendimethalin at 1.0 kg ha^{-1}+hand weeding and hoeing 5 WAE, s-metolachlor at 1.0 kg ha^{-1}+pendimethalin at 1.0 kg ha^{-1}, s-metolachlor at 1.0 kg ha^{-1}+pendimethalin at 0.75 kg ha^{-1}, s-metolachlor at 0.75 kg ha^{-1}+pendimethalin at 1.0 kg ha^{-1}, hand weeding and hoeing 3 WAE, and weedy check). e treatments were laid out in factorial combination ian a

considered as borders. From the end-point of each row, three plants in plots having 10 cm intra row spacing and two plants in 15 cm intra row spacing were considered as borders. us the net plot size was 1.8 m × 1.8 m (3.24 m²). All the recommended practices, except the treatments, were followed to raise the crop. e crop was harvested on 29 October and 6 November 2014 at Sirinka and Jari, respectively. e harvested produce was sun-dried for 7-10 days and threshing and winnowing was done subsequently.

Data collection and analyses

Weeds data: Weed aboveground dry biomass (g m⁻²): For aboveground weed dry biomass, the weeds falling within the quadrate were cut near the soil surface immediately a er recording data on weed count and placed into paper bags separately treatment-wise. e samples were sun-dried for 3-4 days and therea er were placed into an oven at 65°C temperatures till a constant weight and, subsequently, their dry weight was measured. e dry weight was expressed in g m⁻².

Weed Control E ciency (WCE): It was calculated using the following formula:

$$WCE = \frac{(WDC - WDT)}{WDC} \times 100$$

Where,

WDC=Weed dry weight in weedy check, WDT=Weed dry weight in a particular treatment.

Crop data:

Plant height (cm): It was taken with a ruler from 10 randomly taken and pre tagged plants in each net plot area from the base to the apex of the main stem at physiological maturity.

Number of pods per plant: It was taken from the total pods of the above tagged plants at harvest.

Number of per pod: e total number of seeds from the above pods was taken and counted to average the number of seeds pod⁻¹.

Hundred seed weight (g): Out of seeds from the above pods, 100 seeds were counted and their weight was recorded at 10.5% moisture content for hundred seed weight.

Aboveground biomass (g): is parameter was determined by harvesting ten plants in each plot at physiological maturity and their dried aboveground biomass was recorded. Treatment-wise per plant dry weight of straw was multiplied by the number of plants in respective treatments. is was considered as the aboveground dry biomass weight.

Grain yield (kg ha⁻¹): e grain yield was measured a er threshing the sun-dried plants harvested from each net plot and the yield was adjusted at 10.5% seed moisture content. e grain weight obtained in ten plants was added to the nal yield.

Harvest index (%): is parameter was calculated by dividing the grain yield by the aboveground biomass yield and multiplied by 100.

Data analyses: Data on weed density, weed dry biomass, growth, yield attributes and yield were subjected to analysis of variance (ANOVA) using GenStat 15.0 computer so ware [17]. Fisher's protected least signi cant di erence (LSD) test at p 0.05 was used to separate di erences among treatment means [18]. As the F-test of the error variances for most parameters of the two sites was homogeneous combined analysis of data was used.

pigeon pea, effective weed control has been reported with integrated use of pendimethalin and hand weeding [24]. However, lower performance of intra-group herbicides might be due to lower doses than their recommended doses, which needs to be investigated for recommended doses of individual herbicides in mixture [25].

The location and weed management practices interaction further showed that the maximum (472.4 g m^{-2}) mean weed dry weight obtained in weedy check at Jari was significantly higher than all the other interactions (Table 3). These results are consistent with the findings of Arif and Marwat [26,27] who reported more weed dry biomass in weedy check than pre-emergence herbicides (s-metolachlor and pendimethalin) application in canola (*Brassica napus* L.) for weed management.

It was also recognized that there was no significant difference in weed dry weight obtained in weedy check at Sirinka with s-metolachlor at 2.0 kg ha^{-1}, s-metolachlor at 1.0 kg ha^{-1} + pendimethalin at 1.0 kg ha^{-1}, s-metolachlor at 1.0 kg ha^{-1} + pendimethalin at 0.75 kg ha^{-1} and hand weeding at 3 WAE at Jari, and s-metolachlor at 1.0 kg ha^{-1} + pendimethalin at 0.75 kg ha^{-1} and s-metolachlor at 0.75 kg ha^{-1} + pendimethalin at 1.0 kg ha^{-1} at Sirinka. The moderate increase in weed dry weight could be attributed to frequent reoccurrence and persistent characteristics of weeds. Furthermore, like weed density, the weed dry biomass was also lower at Sirinka than at Jari. Herbicide molecules tend to bind with soil clay and organic matter particles, and thus become unavailable for weed killing purposes.

The lower weed dry matter accumulation may be attributed to lower weed density at Sirinka than at Jari (Table 2). The lowest (29.8 g m^{-2}) weed dry weight was recorded with 60 cm × 10 cm planting pattern when treated with s-metolachlor at 1.0 kg ha^{-1} + hand weeding and hoeing 5 WAE, which was statistically at parity with pendimethalin at 1.0 kg ha^{-1} + hand weeding at 5 WAE, s-metolachlor at 1.0 kg ha^{-1} + pendimethalin at 1.0 kg ha^{-1} under both the planting patterns. Jafari [28] stated that pre-emergence herbicides reduced the weed density and dry weight significantly as compared to weedy check in common bean. Similarly, Masoumeh [29] found application of pendimethalin 0.5 kg ha^{-1} + hand weeding 30 days after sowing though was comparable with other treatments, but gave lower weed dry weight after weed-free check in soybean.

The highest (327.5 g m^{-2}) weed dry weight was found in weedy check under 45 cm × 15 cm planting pattern, followed by 60 cm × 10 cm planting pattern and both these interactions resulted in higher increase in weed dry weight than all the other planting pattern and weed management practices interactions (Table 2). A high weed density recorded in the weedy plots invariably resulted in high weed dry weight that could be attributed to low ground cover of cowpea vines.

eldpea with application of s-metolachlor, while the minimum number of seed per pod was obtained in the weedy check plots. Also, the size of pods increased with application of s-metolachlor and hence resulted in maximum number of seeds per pod and vice versa.

Hundred seed weight

Hundred seed weight at Sirinka was 2.4% higher than the hundred seed weight at Jari. e highest (14.0 g) 100 seeds weight was recorded with the application of s-metolachlor at 1.0 kg ha^{-1} + hand weeding and hoeing 5 WAE, which was signi cantly higher (6.4 - 24.3%) than the other treatments (Table 6). e application

Citation: Mekonnen G, Sharma JJ, Negatu LW, Tana T (2016) Growth and Yield Response of Cowpea (Vigna unguiculata L. Walp.) to Integrated Use of Planting Pattern and Herbicide Mixtures in Wollo, Northern Ethiopia. Adv Crop Sci Tech 4: 245. doi: 10.4172/2329-8863.1000245

higher number of seeds per pod and hundred seed weight than that of the control plots. Weedy check had signi cantly lower seed weight and number of seeds per pod than the other treatments.

Aboveground dry biomass weight and grain yield and harvest index were signi cantly higher at Sirinka than at Jari. Application of s-metolachlor at 1.0 kg ha-1+hand weeding and hoeing 5 WAE gave signi cantly higher grain yield and harvest index than other treatments, while it did not signi cantly di er from treatment with pendimethalin at 1.0 kg ha^{-1} supplemented with hand weeding for the aboveground dry biomass yield.

From the result of the study, it can be concluded that managing the weeds with the application of s- metolachlor at 1.0 kg ha^{-1} and weeding and hoeing at 5 WAE along with 60 cm × 10 cm planting pattern proved to be the most pro table practice. Based upon availability, alternatively pendimethalin could also be used in supplement with hand weeding at 5 WAE. Further, to prevent the weed shi , these two herbicides (s-metolachlor and pendimethalin) should be used as herbicide rotation. In future, there is a need to explore the e ectiveness of various combinations of these two herbicides for cost e ective and broad spectrum weed control in cowpea production.

Acknowledgements

We are grateful to the Ministry of Education, Federal Democratic Republic RI (WKLRSLD IRU WKH ¿QDQFLDO VXSSRUW DQG 6LULQND $JULFXOWXUDO 5HVHDUFK &HQWHU for providing research facilities and Mizan Tepi University for permitting the corresponding author for PhD study at Haramaya University.

References

1. Timko MP, Singh BB (2008) Cowpea, a multifunctional legume. In: Moore PH, Ming R, (Eds.) Genomics of tropical crop plants, Springer, New York, pp: 227-257.

2. Pottorff M, Ehlers JD, Fatokun C, Philip A, Close RTJ (2012) Leaf morphology in Cowpea [Vigna unguiculata (L.) Walp]: QTL analysis, physical mapping and identifying a candidate gene using synteny with model legume species. BMC Genomics 13: 234.

3. Diouf D (2011) Recent advances in cowpea [Vigna unguiculata (L.) Walp.]