

# Helicobacter Pylori Infection and Gallbladder Diseases: a Mini-review


Mohammad Javad Yavari Barhaghtalaba<sup>\*1</sup> Mohammad Bagher Jahantaba,<sup>2</sup> Amir Abbas Safaripoura,<sup>2</sup> Reza Hosseinpoura,<sup>2</sup> Saadat Mehrabi Sisakhta, Cambyz Irajieb<sup>2</sup>

FÖ^ælc { ^}ol [-iÖ^a} ^/æ|Ü " ; \*^/ ÉYæ• "W}åç^/•ç^/[-i T^æ&&æ|Ü&^} &^•ÉYæ• "EÖ/æ}

GÖ^ælc { ^}ol [-i T^æ&&æ|Ö{ç^@} ||| \* ÉU@iæ:W}åç^/•ç^/[-i T^æ&&æ|Ü&^} &^•ÉU@iæ:ÉÖ/æ}

**\*Corresponding author:** Mohammad Javad Yavari Barhaghtalaba, Department of General Surgery, Yasuj University of Medical Sciences, Yasuj, Iran, Email-mj.yavari.barhaghtalab@gmail.com

BahiM:ipaloo m



### **Gallbladder cancer**

Risk factors for the development of gallbladder cancer are patient demographics, gallbladder abnormality, patient exposure, and *Salmonella* and *Helicobacter* infections. The association between *H*[21 22]. *pylori* infection and biliary tract carcinoma (gallbladder cancer,