Journal of
Bioremediation & Biodegradation

Jiaet al., J Bioremediat Biodegrad 2017, 8:5
DOI: 10.4172/2155-6199.1000004

Research Article OMICS International

Human Health Risk Levels and Distribution of PAHs in Soil Environment of a Coking Chemical Plant Located in Beijing, China

Jianli Jia [1]*, Haibin He [1], Zihao Wang [1], Shenwei Zhao [1], Linying Me [1], Jialin Liu [1], Shihai Zhang [1], Cunzhen Liang [2] and Yinghua Zhang [3]

[1]School of Chemical and Environmental Engineering, China University of Mining and Technology (Beijing), Beijing 100083, PR China
[2]Beijing Institute of Petrochemical Technology, Beijing 102671, PR China
[3]Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100864, PR China

Abstract

In this study, the polycyclic aromatic hydrocarbons (PAHs) organic pollutants in soil samples were determined by Soxhlet extraction- GCMS methods. Human health risks of as for 5 different sampling sites in the coking chemical plant were evaluated in four major Exposure routes, including oral intake, skin contact, inhalation the air pollutants of outdoor which from the soil surface, and inhalation the air pollutants of outdoor which from the soil lower layer. The results showed that the detection rate of 16 kinds precedent- controlled PAHs in the US Environmental Protection Agency

the oral exposure pathway is 498 times the carcinogenic risk of the skin exposure pathway, and 198 times the rate of carcinogenic risk of inhalation of soil particle Exposure route. And the rate of carcinogenic risk of Inhalation the air pollutants of outdoor which from the soil lower layer is almost 0. The hazard quotient of soil ∉PAHs main air pollutants pathway are oral intake and inhalation of outdoor air from the soil surface. The sum of them is close to 90%, The hazard quotient of soil ∉PAHs in the coking plant area is seriously exceeded. The risk of ∉PAHs in the soil of coking plant was mainly exposed to oral intake, and the contribution rate of carcinogenic risk was 99.3103%. Combined with the development characteristics of China's coking industry, regional climate and geological characteristics, make a human health risk assessment to ∉PAHs.

Keywords: Carcinogenic risk; Hazard quotient; Coking industry; PAHs; Soil

Introduction

*Corresponding author: Jianli Jia, School of Chemical and Environmental Engineering, China University of Mining and Technology (Beijing), Beijing 100083, PR China, Tel: 13810103722; E-mail: jiajianli77@126.com

Received July 27, 2017; Accepted August 22, 2017; Published August 24, 2017

Citation: Jia J, He H, Wang Z, Zhao S, Me L, et al. (2017) Human Health Risk Levels and Distribution of PAHs in Soil Environment of a Coking Chemical Plant Located in Beijing, China. J Bioremediat Biodegrad 8: 407. doi: 10.4172/2155-6199.1000407

Copyright: ' 2017 Jia J, et al. This is an open-a ccess article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Formula number	Exposure route	Formula Description	Exposure formula expression
9	Oral intake	Carcinogenic risk	$CR_{ois} = OISER_{ca} \times C_{sur} \times SF_o$
10		Hazard quotient	$HQ_{ois} = \frac{OISER_{nc} \times C_{sur}}{RfD_o \times SAF}$
11	Skin contact	Carcinogenic risk	$CR_{dcs} = DCSER_{ca} \times C_{sur} \times SF_d$
12		Hazard quotient	$HQ_{dcs} = \frac{DCSER_{nc} \times C_{sur}}{RfD_d \times SAF}$
13	Inhalation the air pollutants of outdoor which from the soil surface	Carcinogenic risk	$CR_{iov1} = IOVER_{ca1} \times C_{sur} \times SF_i$
14		Hazard quotient	$HQ_{iov1} = \frac{IOVER_{nc1} \times C_{sur}}{RfD_i \times SAF}$
15	Inhalation the air pollutants of outdoor which from the soil lower layer	Carcinogenic risk	$CR_{iov2} = IOVER_{ca2} \times C_{sub} \times SF_i$
16		Hazard quotient	$HQ_{iov2} = \frac{IOVER_{nc2} \times C_{sub}}{RfD_i \times SAF}$

Table 4: Cancer risk and hazard quotient calculating formulas for four soil exposure pathways.

quotient of inhalation the air pollutants of outdoor which from the soil lower layer, Dimensionless.

Based on total carcinogenic risk level, calculating separately the contribution rate of carcinogenic risk in the four exposure pathways (Formula 1), analyzing and determining the main contribution path to provide the basis for the follow- up risk prevention and control.

$$R_i = \frac{CR_i}{\sum CR_i} \times 100 \quad \text{(Formula 1)}$$

And R_i- e contribution rate of carcinogenic risk or hazard quotient under the exposure pathways, Dimensionless; CR_i Level of carcinogenic risk or hazard quotient under the exposure pathways, Dimensionless. CR_i Total carcinogenic risk or total hazard quotient.

Risk control value calculation: When the level of carcinogenic risk or hazard quotient exceeds the acceptable level, the risk control value for the corresponding exposure pathway should be calculated (Table 5).

Results and Discussion

Characteristics and PAHs content in coking plant

e contents of soil pH, water and PAHs in the sampling sites of coking plant are shown in Table 6. e results showed that soil in coking plant is basically weak acidity, pH distribution between 6.5-7, soil moisture content is generally low, the moisture content of 14 samples are between 10% and 20%, and 15% of the total sample are less than 5%, e PAHs content in the soil of the coking plant was 0.88-447.24 mg/kg, e content of Baa at 100 m was 447.24 mg/kg, and the content of Bbf at 100 m was 380.29 mg/kg, ere is little di erence between 0 m sampling and 200 m sampling of PAHs content, but the PAHs content at 100 m was the largest.

Human health risk assessment of soil in coking plant

Carcinogenic risk: According to Tables 1-4 evaluation methods and the main sk asi5anment of soT-10.9on metf.5 (o)12 (f)0.6 712 66 (t i)3

Sampling point (m)	pH	Moisture content%	types of pollutant	Average content (mg/kg)	Oral intake		Skin contact		Inhalation the air pollutants of outdoor which from the soil surface		Inhalation the air pollutants of outdoor which from the soil lower layer	
					Carcinogenic risk	Hazard quotient	Carcinogenic risk	Hazard quotient	Carcinogenic risk	Hazard quotiet	Carcinogenic risk	Hazard quotient
0 m	6.7	15.51	×PAHs	164.80	5.88E-03	7.94	1.18E-05	2.53	2.42E-05	9.37	4.80E-09	1.33E-03
			Baa	25.06	8.94E-04	1.21	1.80E-06	0.38	2.56E-06	0.99	1.66E-11	4.58E-06
			Bbf	27.68	9.87E-04	1.33	1.98E-06	0.42	4.26E-07	0.17	4.15E-13	1.15E-07
			Bap	8.48	3.03E-04	0.41	6.08E-07	0.13	1.18E-07	0.05	1.04E-13	2.88E-08
50 m	6.8	13.33	×PAHs	599.07	2.14E-02	28.86	4.30E-05	9.20	1.26E-04	48.87	3.61E-08	9.97E-03
			Baa	45.15	1.61E-03	2.17	3.24E-06	0.69	4.61E-06	1.79	2.98E-11	8.25E-06
			Bbf	8.21	2.93E-04	0.40	5.89E-07	0.13	1.26E-07	0.05	1.23E-13	3.40E-08
			Bap	11.45	4.09E-04	0.55	8.21E-07	0.18	1.59E-07	0.06	1.41E-13	3.89E-08
100 m	7	10.14	×PAHs	1378.9	4.92E-02	66.42	9.89E-05	21.17	1.88E-04	73.04	6.96E-08	1.92E-02
			Baa	447.24	1.60E-02	21.54	3.21E-05	6.87	4.57E-05	17.71	2.96E-10	8.17E-05
			Bbf	380.29	1.36E-02	18.32	2.73E-05	5.84	5.85E-06	2.27	5.70E-12	1.58E-06
			Bap	91.49	3.26E-03	4.41	6.56E-06	1.40	1.27E-06	0.49	1.12E-12	3.10E-07
150 m	6.9	11.57	×PAHs	923.79	3.30E-02	44.50	6.62E-05	14.18	1.85E-04	71.80	3.63E-08	1.00E-02
			Baa	104.48	3.73E-03	5.03	7.49E-06	1.60	1.07E-05	4.14	6.91E-11	1.91E-05
			Bbf	112.09	4.00E-03	5.40	8.04E-06	1.72	1.72E-06	0.67	1.68E-12	4.64E-07
			Bap	58.02	2.07E-03	2.79	4.16E-06	0.89	8.08E-07	0.31	7.13E-13	1.97E-07
200 m	6.5	5.75	×PAHs	204.11	7.28E-03	9.83	1.46E-05	3.13	6.31E-05	24.48	3.29E-08	9.10E-03
			Baa	10.71	3.82E-04	0.52	7.68E-07	0.16	1.09E-06	0.42	7.08E-12	1.96E-06
			Bbf	20.62	7.35E-04	0.99	1.48E-06	0.32	3.17E-07	0.12	3.09E-13	8.54E-08
			Bap	14.39	5.13E-04	0.69	1.03E-06	0.22	2.00E-07	0.08	1.77E-13	4.88E-08

Table 6: Sample test and evaluation results.



Figure 4: Hazard quotient of different sampling sites in the coking plant.

(a) Carcinogenic risk.

(b) hazard quotient.

Figure 5: The contribution ratio of different exposure pathways to human risk.

(a) Carcinogenic risk contribution rate of major pollutants

(b) Hazard quotient contribution rate of major pollutants

Figure 6: The proportion of the main pollutants in the soil of coking plant to total pollutants.

Citation: Jia J, He H, Wang Z, Zhao S, Me L, et al. (2017) Human Health Risk Levels and Distribution of PAHs in Soil Environment of a Coking Chemical Plant Located in Beijing, China. J Bioremediat Biodegrad 8: 407. doi: 10.4172/2155-6199.1000407

5. Wild SR, Jones KC (1995) Polynuclear aromatic hydrocarbons in the United Kingdom environment a preliminary source inventory and budget. Environmental Pollution 88: 91-108.

6. Jones KC, Stratford JA, Waterhouse KS, Furlong ET, Giger W, et al. (1989) Increases in the polynuclear aromatic hydrocarbon content of an agricultural soil over the last century. Environmental Science & Technology 23: 95-101.

7. Kulh§nek A, Trapp S, Sismilich M, JankŢ J, Zimov§ M (2005) Crop-speciýc human exposure assessment for polycyclic aromatic hydrocarbons in Czech soils. Science of the Total Environment 339: 71-80.

8. Jia J, Li X, Yang L, Hu L, Zhao SW, et al. (2016) Soil northwest China region in a coal chemical arsenic risk to human health and the safety threshold. Earth Science Frontiers 23: 124-132.

9. Jia J, Li X, Wu P, Liu Y, Han C, et al. (2015) Human health risk assessment and safety threshold of harmful trace elements in the soil environment of the Wulantuga open-cast coal mine. Minerals 5: 837-848.

10. Liu G, Niu J, Guo W, An X, Zhao L (2016) Ecological and health risk-based characterization of agricultural soils contaminated with polycyclic aromatic hydrocarbons in the vicinity of a chemical plant in China. Chemosphere 163: 461-470.

11. Yi-Zhong W, Xiao Liang T, Ying GE (2011) Comparison of RBCA and Csoil Models in Health Risk Assessment of Volatile Organic Pollutants. Journal of Agricultural Environmental Sciences 30: 2458-2466.

12. Cao Y, Zhang J (2009) Comparison of RBCA and CLEA Models in Environmental Risk Assessment of Heavy Metal Contaminated Sites. Environmental Sciences 22: 241-247.

13. Lemke LD, Bahrou AS (2009) Partitioned multi objective risk modeling of carcinogenic compounds in groundwater. Stochastic Environmental Research and Risk Assessment 23: 27-39.

14. Waitz MFW, Freijer JI, Kreule P, Swartjes FA (1996) The VOLASOIL risk assessment model based on CSOIL for soils contaminated with volatile compounds. RIVM Rapport, pp: 7158-10014.

15. Ming Z, Tian-Tian C, Lin S (2013) CLEA model and its application in China. Journal of University of Chinese Academy of Sciences 30: 779-785.

16. Meng-Fang C, Yong-Ming L, Jing S (2011) Theory and Common Model of Soil General Assessment Basis for Contaminated Sites. Environmental Monitoring Management and Technology 23: 19-25.

17. National Environmental Protection Standard of the People's Republic of China (2014) Technical Guidelines for Risk Assessment of Contaminated Sites.

18. You-Ning X, Jiang-Hua Z, Hai-Ling KE, Hua-Qing C, Gang Q, et al. (2014) Human health risk of heavy metal pollution in farmland soil in a gold mining area. Journal of Geology and Mineral Resources 8: 1239-1252.

19. IARC (2006) Polycyclic Aromatic Hydrocarbons. IARC Monographs, Lyone, France.

20. Singh