



Impact of Agroecological Practices on Soil Organic Carbon Stocks in the Central Highlands of Madagascar: Case of Itasy Region

Narindra Rakotovao

Laboratoire des Radiosotopes, Université d'Antananarivo, Route d'Andraisoro B.P. 3383, Antananarivo 101, Madagascar.

Abstract:

In the Itasy Region of Madagascar, agroecological practices were promoted to improve farmers' income while ensuring sustainable management of natural resources. These practices were agroforestry (AF), on-farm tree plantation (TP) and the use of organic fertilizers such as compost and improved manure (CIM). This study aimed to assess the impact of each of these agroecological practices on soil organic carbon (SOC) stock versus conventional practices. Soil survey was carried out on 712 farmers' fields in order to know the practices' history of each plot and to define chronosequences expressing the range of agroecological practice duration. Composite soil samples at three depths (0-10, 10-20 and 20-30 cm) were collected from these 712 farmers' plots to evaluate SOC stocks; the two main soil types of the Itasy Region are Ferralsols and Andosols. Results showed that soil type was a main determinant of the spatial variability of SOC. Agroforestry based on fruit species associated with annual cash crops (legume, tomato) aging more than 8 years and fertilized regularly with CIM at a rate of 8.4 t ha⁻¹ presented the highest SOC sequestration rate estimated at 2.4 MgC ha⁻¹ yr⁻¹ compared to annual crops without fertilization. SOC sequestration rate of 0.4 MgC ha⁻¹ yr⁻¹ was observed on growing vegetables for market fertilized with CIM at a rate of 10.2 t ha⁻¹. SOC stocks for TP sites presented no



difference compared to traditional practices, because the

Citation: Narindra Rakotovao; Impact of Agroecological Practices on Soil Organic Carbon Stocks in the Central Highlands of Madagascar: Case of Itasy Region; Agri 2020; April 21, 2020; Venice, Italy