

Improving Global Assessment of Functioning (GAF)

Monrad Aas I. H*

Research Unit, Division of Mental Health and Addiction, Vestfold Hospital Trust, 3103 Tönsberg, Norway

SUMMARY

A large number of scoring systems are developed for psychiatry. The Global Assessment of Functioning (GAF) is known worldwide, translated into many languages, and used in many outcome studies. GAF is used to rate severity of illness in psychiatry and covers the range from positive mental health to severe psychopathology. It is an overall (global) measure of how patients are doing. GAF is not intended to be a diagnosis-specific scoring system, but a generic. Compared to diagnosis, GAF values represent more multidimensional information. With GAF, the degree of mental illness is measured by rating psychological, social and occupational functioning [Aas, 2010; 2011; 2014a].

From research, we know there are problems with GAF (for example, reliability and validity problems). The properties of the GAF scale need a closer examination with the potential for improvement in mind. The present study was published in *International Journal of Emergency Mental Health* [2014b] and has focus on GAF properties. Purpose: to show both gaps in current knowledge and ideas for further development. Methods: The study is based upon a systematic literature review. Findings: for the properties of GAF, numerous gaps in knowledge were found: for example, a continuous scale is used for the present GAF, but would a categorical scale make a better GAF? On visual scales scoring is done by setting a mark directly on the scale, but would transformation to a visual scale result in an improved GAF? The anchor points (including examples) were decided early in the history of GAF, but would new anchor points and examples result in a better GAF (anchor points for symptoms, functioning, positive mental health, prognosis, improvement of

generic properties, exclusion criteria for scoring in each 10-point intervals, and anchor points at the endpoints of the scale)? Is a change in the number of anchor points and their distribution over the total scale important? Rating within 10-point intervals can be requiring, but can better instructions improve this? Internationally, GAF with both one and two values are used, but what is the advantage of having separate symptom (GAF-S) and functioning scales (GAF-F)? GAF-S and GAF-F scales should score different dimensions and still be correlated, but what is the best combination of definitions for GAF-S and GAF-F? Conclusions: Given the widespread use, research-based development of GAF has not been especially strong. Further research could improve GAF.

REFERENCES

- Aas, I.H.M. (2010). Global Assessment of Functioning (GAF): Properties and frontier of current knowledge. *International Journal of Emergency Mental Health and Human Resilience*, *10*, 1-2.
- Aas, I.H.M. (2014a). Collecting information for rating Global Assessment of Functioning (GAF): Sources of Information and Methods for Information Collection. *Current Psychiatry Reviews*, *10*(4), 330-347.
- Aas, I.H.M. (2014b). Towards a better Global Assessment of Functioning (GAF): improving scale properties. *International Journal of Emergency Mental Health and Human Resilience*, *17*, 197-202.

*Correspondence regarding this article should be directed to: monrad.aas@siv.no; monradaas@yahoo.no