
*Corresponding author: Francesca Milano, Private Practice of Orthodontics and Dental Sleep Medicine, Bologna, via Clavature 1, 40124 Bologna, Italy, Tel: +39 051 228084; Fax: +39 051 239889; E-mail: francescamilano@libero.it

Received November 18, 2013; Accepted December 07, 2013; Published December 09, 2013

Citation: Milano F (2013) The Value of a Multidisciplinary Integrated Approach

- ciation of hypertension and sleep-disordered breathing. Arch Intern Med 160: 2289-2295.
2. Budhiraja R, Parthasarathy S, Quan SF (2007) Endothelial dysfunction in obstructive sleep apnea. J Clin Sleep Med 3: 409-415.
 3. Gottlieb DJ, Yenokyan G, Newman AB, O'Connor GT, Punjabi NM, et al. (2010) Prospective study of obstructive sleep apnea and incident coronary heart disease and heart failure: the sleep heart health study. Circulation 122: 352-360.
 4. Kario K (2009) Obstructive sleep apnea syndrome and hypertension: ambulatory blood pressure. Hypertens Res 32: 428-432.
 5. Neau JP, Paquereau J, Meurice JC, Chavagnat JJ, Gil R (2002) Stroke and sleep apnoea: cause or consequence? Sleep Med Rev 6: 457-469.
 6. Pamidi S, Tasali E (2012) Obstructive sleep apnea and type 2 diabetes: is there a link? Front Neurol 3: 126.
 7. Wang X, Bi Y, Zhang Q, Pan F (2013) Obstructive sleep apnoea and the risk of type 2 diabetes: a meta-analysis of prospective cohort studies. Respirology 18: 140-146.
 8. Guilleminault C, Stoohs R, Clerk A, Cetel M, Maistros P (1993) A cause of excessive daytime sleepiness. The upper airway resistance syndrome. Chest 104: 781-787.
 - 9.