

**Open Access** 

 $\begin{array}{c} \mathbf{K} & := - \left( l \right) = \left( \frac{1}{2} \right) \left( \frac{1}{2} \right)$ 

Citation: Yahou X (2023) In common Wheat, Yield-related Traits are Enhanced by the Introduction of QTL from Aegilops Tauschii. J Plant Genet Breed 7: 152.

í =-1.44 . 10%, 1 . ľ K -14 ď \* τ -T 1 1. Ш. - 1 ł J ; ( y 😃 015 J ... **₹**-† Ł ÷ 1 .... 1. ł ٠t = -10.

**،** ۹ 1<u>5</u> "; t ! 1 - l ~ 14 ..... 3 -24 J g ię. ę 1 ÷ 1 ÷ę 2 .... ł ł ١ 0 4). ( t\_ . . . ! Į -114 N.

015/ 1 L a a = -1 anat ۰., 55 15. 131 1 I\_ ¥ ď **ş** 1 1 ¢ ₽4<sup>6</sup> 9 <sup>6</sup> 9₽4 • • • • • • • • • • 506 1426 ę v . J 1 -. 41 1 t • 12 ٩y X С

55 ÷ 1 -44 ۰**v** 44 . . . 15 ÷ŧ ÷ .1.4 61-0 20 1 015. ſ ÷ • 015 J. 1.1 ł 2 Ø 1\_100 00.1. 32 ł 6 1 1. 015 .1 . 1 • ] • ۹, je ۶*۲* ',• \ ŗ ÷ 1 -14 Q. \* 19.90 T ₽, 1 . J at your ď J Curl \* " · . . . 1 74 А 1 J •-• t v

## A⊠

C 🛛 I

## References

- Mcfadden ES, Seal ER (1946) The origin of Triticum spelta and its freethreshing hexaploid relatives. J Hered 37: 81-89.
- K[ \* H, Zhang Z, Yang Y, Wei C, X\* L, et al. (2023) Advances in the Mining of Disease Resistance Genes from Aegilops tauschii and the Utilization in Wheat. Plants (Basel) 12: 880.
- H<sup>\*</sup>ang S, Siłikhach [Inkic A, S<sup>\*</sup> X, Fali• J, Gill B, et al. (2002) Genes encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the Triticum/ Aegilops complex and the evolutionary history of polyploid wheat. Proc Natl Acad Sci U S A 99: 8133-8138.
- 4. Błaglia L, Gaça::i F, M[!ell[ L, et al. (2020) On the applicability of the Tubulin-Based Polymorphism (TBP) genotyping method; comprehensive guide ill\*•claced ch![`gh che a]]licaci[n [n di elent genetic le line• in che leg`me family