Influence of Single Nucleotide Polymorphisms on the Efficacy of Platelet-Rich Plasma in the Treatment of Injuries

Ricard Pruna*

F.C. Barcelona Medical Head Services, FIFA Medical Center of Excellence, Barcelona, Spain

***Corresponding author:** Ricard Pruna, F.C. Barcelona Medical Head Services, FIFA Medical Center of Excellence, Barcelona, Spain, E-mail: ricard.pruna@barcelona.cat

Received date: January 04, 2016; **Accepted date:** January 25, 2016; **Published date:** February 17, 2016

Copyright: © 2016 Pruna R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Single Nucleotide Polymorphisms in Platelet-Rich Plasma

Tissue repair in musculoskeletal injuries is often a slow and sometimes incomplete process. Musculoskeletal injuries have a great impact on athletes' especially elite athletes – and a rapid recovery of full efficiency and return to competition is of primary importance. The search for a minimally invasive treatment of these injuries is of great importance, especially in the world of sports.

The use of growth factors is thought to be useful in clinical practice because it promotes rapid healing with a high-quality tissue and allows a nearly and safe return to unrestricted activity. Platelet-rich plasma (PRP) is a simple and minimally invasive way to obtain a natural concentration of autologous growth factors, including IGF, EGF, TGFb1, and FGF2.

PRP is currently being widely experimented in