Epidemiology: Open Access

Debottam et al., Epidemiology (Sunnyvale) 2016, 6:2
10.4172/2161-1165.1000228

Malnutrition Scenario among School Children in Eastern-India-an Epidemiological Study

Debottam Pal[1], **Suman Kanungo**[1], **Baishali Bal**[1], **Kalyan Bhowmik**[1], **Tanmay Mahapatra**[1] **and Kamalesh Sarkar**[1,2*]

[1]*National Institute of Cholera and Enteric Diseases, Kolkata, India*

[2]*National Nutrition Monitoring Bureau, West Bengal Unit, Kolkata, India*

[*]**Corresponding author:** Kamalesh Sarkar, Scientist F, National Institute of Cholera & Enteric Diseases and Officer-in-Charge, National Nutrition Monitoring Bureau, West Bengal Unit, P-33, CIT Road, Scheme - XM, Beliaghata, India, Tel: +91-33-23631222; E-mail: kamalesh.sarkar@gmail.com

Received date: February 19, 2016; **Accepted date**: March 03, 2016; **Published date**: March 10, 2016

Copyright: É 2016 Sarkar K, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Twenty years follow after the implementation (1995) of the school-meal program to improve nutritional status, drop out from school and academic efficiency among school-children, malnutrition remained a major concern in India. Efforts to quantify the problem and identify its determinants were handful, especially in the eastern part of the country. Thus the purpose of this study was to determine the burden and possible predictors of malnutrition among primary and upper-primary school-children in eastern India.

Methods: Using stratified cluster random sampling method, a multistage cross-sectional study was conducted involving all twenty educational districts of West Bengal, a highly populous state in eastern India. During 2014-15, using structured questionnaire, standard anthropometry and laboratory testing a representative sample of 24,108 primary and upper-primary students from the whole state of West Bengal were interviewed and assessed. Descriptive and regression analyses were conducted using SAS-9.4.

Results: Among 24,108 recruited students aged between six and thirteen years, prevalence of under-nutrition was alarmingly high (about twenty-three percent). Furthermore, over half of the students (fifty-four percent) were at risk of developing under-nutrition. On the other hand, only seventeen percent of the primary students had ideal nutritional status. Odds of being malnourished were higher among male students (compared to females), those belonging to younger age (studying in primary compared to upper-primary classes), Muslim religion (with reference to Hindus), and under- privileged caste (in comparison with general caste) as well as those residing in rural areas (as opposed to urban). Parental education was negatively associated with the likelihood of under-nutrition. Those who had more than three siblings, unemployed father and students with maternal death were more likely to be undernourished.

Conclusions: Prevalence of under-nutrition was high among school-children in the study area. School-based interventions targeting high-risk, under-privileged children, especially in rural areas with lower parental education and poor level of sanitary practices seemed to be urgently required.

Keywords Adolescent nutrition; Child nutrition, Malnutrition; Nutritional epidemiology; Indian children; Determinants of malnutrition

Introduction

Brain development of a child starts in mother's womb and continues throughout infancy and up to the age of 2 year, whereas its growth continues beyond that period. Appropriate intake of macro and micro-nutrients during intra-uterine life as well as few years following birth are crucial for brain and other organ development [1]. Under-nutrition during this period can slow down the physical and mental development of the child and consequences are usually lifelong [2,3].

Poor socio-economic conditions in developing and under-developed countries result in under-nutrition since intrauterine life till early childhood and attempts must be made to ensure their appropriate nutrition during school-age. Under-nutrition and infections like diarrhoea, respiratory infections, malaria, TB etc. are inter-dependent and parts of a vicious cycle [4-6].

On the other hand, chronic infections make children undernourished [5,7]. Prolonged under-nutrition has detrimental on all organ-systems including gut, reducing its capacity to maintain homeostasis during under-nutrition and may result into life threatening conditions [8-11].

Dietary requirement of essential nutrients varies with age, gender, physiological status and physical activity [12]. Globally, millions of school-going children from under-nutrition [13]. In India, National Family Health Survey-3 (NFHS-3) estimated that in 2006, 24% under-5 years-aged children were severely malnourished (<3 standard deviation of the reference value suggested by the Indian Academy of Paediatrics) according to the height-for-age scale and 16% according to the weight-for-age values.

Free and compulsory education till the age of 14 year is constitutional right, and institutional commitment in India. Despite these measures, about 40% children drop out during their primary schooling [14].