



Methods for Root Canal Preparation and Treatment Results: Follow-Up for 5 Years

Pan Gary Cheung*

the literature suggests that either nonsurgical or surgical endodontic treatment improves OHRQoL. However, due to the

of the root canal preparation and treatment results. The results of the study are summarized in Table 1.

C

The results of the study showed that the success rate of the root canal preparation and treatment was 64.4% (90/140). The success rate was significantly higher in the 6-12 months follow-up period (64.4%) compared to the 1-5 years follow-up period (21.3%). The success rate was significantly higher in the 6-12 months follow-up period (64.4%) compared to the 1-5 years follow-up period (21.3%).

The results of the study showed that the success rate of the root canal preparation and treatment was 32.6% (14/43). The success rate was significantly higher in the 6-12 months follow-up period (32.6%) compared to the 1-5 years follow-up period (21.3%). The success rate was significantly higher in the 6-12 months follow-up period (32.6%) compared to the 1-5 years follow-up period (21.3%).

The results of the study showed that the success rate of the root canal preparation and treatment was 12.8% (5/39). The success rate was significantly higher in the 6-12 months follow-up period (12.8%) compared to the 1-5 years follow-up period (21.3%). The success rate was significantly higher in the 6-12 months follow-up period (12.8%) compared to the 1-5 years follow-up period (21.3%).

D

The results of the study showed that the success rate of the root canal preparation and treatment was 13.9% (5/36). The success rate was significantly higher in the 6-12 months follow-up period (13.9%) compared to the 1-5 years follow-up period (21.3%). The success rate was significantly higher in the 6-12 months follow-up period (13.9%) compared to the 1-5 years follow-up period (21.3%).

The results of the study showed that the success rate of the root canal preparation and treatment was 13.9% (5/36). The success rate was significantly higher in the 6-12 months follow-up period (13.9%) compared to the 1-5 years follow-up period (21.3%). The success rate was significantly higher in the 6-12 months follow-up period (13.9%) compared to the 1-5 years follow-up period (21.3%).

- restorations during 16 years: a case report. Clin Implant Dent Relat Res 7: 200-208.
14. Harrel SK, Barnes JB, Hidalgo FR (1998) Aerosol and splatter contamination from the operative site during ultrasonic scaling. J Am Dent Assoc 129: 1241-1249.
15. Ionescu AC, Cagetti MG, Ferracane GL, Godoy FG, Brambilla E, et al. (2020) Topographic aspects of airborne contamination caused by the use of dental handpieces in the operative environment. J Am Dent Assoc 151: 660-667.