# Migraine and Lactose Malabsorption: Is there Any Relationship?

**Del Zompo F, Mangiola F, Petruzziello C, D'Angelo G, Di Rienzo TA, Franceschi F, Gasbarrini A and Ojetti V***

*Department of Internal Medicine and Gastroenterology, Catholic University of the Sacred Heart, Rome, Italy*

***Corresponding author:** Veronica Ojetti, Department of Internal medicine and Gastroenterology, Catholic University of the Sacred Heart, Rome, Italy, Tel: 00390630153188; E-mail:

**Rec date:** Nov 18, 2015, **Acc date:** Jan 21, 2016, **Pub date:** Jan 28, 2016

**Copyright:** © 2016 [illegible] DQG VHYHULW\ RI KHDGDFKH GXULQJ WKH /%7 DQG D FRUUHODWLRQ ZLWK /0

**Patients and Methods:** We enrolled 93 (74F/19M; mean-age 42 ± 16 yrs) outpatients who performed a LBT according to the guidelines. Patients reported on a Visual-Analogue-Scale the intensity of headache before and at the end of the test. We considered as significant an increase of headache in VAS peak >10mm.

**Results:** 67/93 (72%) resulted LM, while 53% (49/93) referred a headache worsening. Within the group of LM headache worsened in 50.7%, compared to 57.7% in non-malabsorbers (p=0.36). In the subgroup of patients with habitual headaches there is a significant worsening compared to non-habitual ones (70.7% vs 38.5%; p=0.003) regardless of the LBT result.

**Conclusion:** The present study denies the existence of an association between LM and headache. We found that headache significantly worsened in patients usually suffering from headache, regardless LM. We think that this data should be considered, since we suppose that the onset of headache during a LBT could affect the correct execution of the test.