

# MRI Reveals Comorbid Cerebrovascular Lesions in Glaucoma

Christopher Lai\*

*The Hong Kong Polytechnic University, Hung Hom, HKSAR, China*

\*Corresponding author: Department of Health Technology and Informatics, The Hong Kong Polytechnic University, Hung Hom, HKSAR, China, Tel: +852-3400-8596; Fax: +852-2362-4365; E-mail: chris.lai@polyu.edu.hk

Received date: April 15, 2015, Accepted date: April 16, 2015, Published date: April 19, 2015

Copyright: © 2015 Lai C, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Editorial

Everyone above the age of 40 years is at risk for glaucoma, particularly in those with diabetes, a familial history of glaucoma and a history of eye injury. It is one of the dominating causes of blindness in worldwide. Owing to its asymptomatic nature, a significant, irreversible and progressive centripetal visual field loss subsequent to the damage of the optic nerve head and the loss of retinal nerve fiber layer are commonly found in newly diagnosed individuals. Today, the main modifiable treatment option to delay the progression of glaucoma and preserve the visual field remains relying on the reduction of intraocular pressure. However, rigorous control of

ip°  
° If r ° ° q of  
g ap inf Mq eg inM° of r M° —M g n sa lB M

---

ischemic changes in cerebral magnetic resonance imaging. Jpn J Ophthalmol 48: 340-344.

10. Kitsos G, Zikou AK, Bagli E, Kosta P, Argyropoulou MI (2009) Conventional MRI and magnetisation transfer imaging of the brain and optic pathway in primary open-angle glaucoma. Br J Radiol 82: 896-900.