

# Neurodevelopment, Intestinal Function, and Autism

Edwin Chau-Leung Yu\*

*Intemed Hong Kong, Shan Tung Street, Mongkok, Kowloon, Hong Kong*

\*Corresponding author: Edwin Chau-leung Yu, Intemed Hong Kong, Shan Tung Street, Mongkok, Kowloon, Hong Kong, Tel: +85223012418; E-mail: [yuedwin@yahoo.com](mailto:yuedwin@yahoo.com)

Received Date: Aug 20, 2018; Accepted Date: Sep 14, 2018; Published Date: Sep 22, 2018

Copyright: © 2018 Yu ECL. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use,

## Gut Influence on Brain Development

Visceral sensations develop as the enteric nervous system (ENS),

acceptance improved mostly in that sequence (Figure 1). 5 had treatment stopped within 1.5-2 years, while one with Asperger syndrome continued with stubbornness recovered after 25 years though negative and grudging thoughts changed positive only after 3 years. One child mainly stubborn was not compliant, feeding improved but worsened with frequent running nose and cough, and overall recovery variable with pattern not distinct. Follow-up in 2018 could not reach this child, being in foreign lands. For the remainder 5 patients, follow-up showed, for school: results good or upper grade in 3

representations are processed and activated by the hippocampus in



- 34 Clarke G, Grenham S, Scully P, Fitzgerald P, Moloney RD, et al. (2013) The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Mol Psychiat* 18: 666-673.
- 35 Mayer EA (2011) Gut feelings: the emerging biology of gut-brain communication. *Nat Rev Neurosci* 12: 453-466.
- 36 Grenham S, Clarke G, Cryan JF, Dinan TG (2011) Brain-gut-microbe communication in health and disease. *Front Physiol* 2: 94.
- 37 Cryan JF, O'Mahony SM (2011) The microbiome-gut-brain axis from bowel to behavior. *Neurogastroenterol Motil* 23: 187-192.
- 38 Diaz Heijtz R, Wang S, Anuar F, Qian Y, Björkholm B, et al. Normal (2011) gut microbiota modulates brain development and behavior. *Proc Natl Acad Sci USA* 108: 3047-3052.
- 39 Hsiao EY, McBride SW, Hsien S, Sharon G, Hyde ER, et al. (2013) Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders. *Cell* 155: 1451-1463.
- 40 De Palma G, Blennerhassett P, Lu J, Deng Y, Park AJ, et al. (2015) Microbiota and host determinants of behavioural phenotype in maternally separated mice. *Nat Commun* 6: 7735.
- 41 Desbonnet

- 80 Inui T, Kumagaya S, Myowa-Yamakoshi M (2017) Neurodevelopmental Hypothesis about the Etiology of Autism Spectrum Disorders. *Frontiers in Human Neuroscience* 11: 354.
81. American Psychiatric Association (2013) Diagnostic and Statistical Manual of Mental Disorders 6th Edn. American Psychiatric Press, Washington DC, US.
82. Zablotsky B, Black LI, Maenner MJ, Schieve LA, Blumberg SJ (2014) Estimated prevalence of autism and other developmental disabilities following questionnaire changes in the 2014 national health interview survey. *Natl Health Stat Report* 13: 1-20.
83. Clements CC, Zoltowski AR, Yankowitz LD, Yerys BE, Schultz RT, et al. (2018) Evaluation of the Social Motivation Hypothesis of AutismA Systematic Review and Meta-analysis. *JAMA Psychiatry* 75: 797-808.
84. Nover A (2018) fMRI Evidence for Social Motivation Hypothesis of Autism. *Psychiatry Advisor*.
85. Scott-Van Zeeland AA, Dapretto M, Ghahremani DG, Poldrack RA, Bookheimer SY (2010) Reward processing in autism. *Autism Res* 3: 53-67.
86. Pavl D (2017) A Dopamine Hypothesis of Autism Spectrum Disorder. *Dev Neurosci* 39: 355-360.
87. McNamara CG, Dupret D (2017) Two sources of dopamine for the hippocampus. *Trends in Neurosciences* 40: 383-384.
88. Cooper RA, Richter FR, Bays PM, Plaisted-Grant KC, Baron-Cohen S, et al. (2017) Reduced Hippocampal Functional Connectivity During Episodic Memory Retrieval in Autism. *Cerebral Cortex* 27: 888-902.
89. Marin-Burgin A, Schinder AF (2012) Requirement of adult-born neurons for hippocampus-dependent learning. *Behav Brain Res* 227: 391-399.
90. Snyder JS, Soumier A, Brewer M, Pickel J, Cameron HA (2011) Adult hippocampal neurogenesis buffers stress responses and depressive behaviour. *Nature* 476: 458-461.
91. Desbonnet L, Clarke G, Shanahan F, Dinan TG, Cryan JF (2014) Microbiota is essential for social development in the mouse. *Mol Psychiatry* 19: 146-148.
92. de Tijae CG, Wöpereis H, Ramadan M, van Eijndthoven T, Knol J, et al. (2014) Altered gut microbiota and activity in a murine model of autism spectrum disorders. *Brain Behav Immun* 37: 197-206.
93. Douglas-Escobar M, Elliott E, Neu J (2013) Effect of intestinal microbial ecology on the developing brain. *JAMA Pediatr* 167: 374-379.
94. Mulle JG, Sharp WG, Cubells JF (2013) The gut microbiome: a new frontier in autism research. *Curr Psychiatry Rep* 15: 337.
95. MacFabe DF (2012) Short-chain fatty acid fermentation products of the gut microbiome: implications in autism spectrum disorders. *Microb Ecol Health Dis* 23: 19260.
96. Doenças C (2018) Gut Microbiota, Inflammation, and Probiotics on Neural Development in Autism Spectrum Disorder. *Neuroscience* 374: 271-286.
97. Iovene MR, Bombace F, Maresca R, Sapone A, Iardino P, et al. (2017) Intestinal Dysbiosis and Yeast Isolation in Stool of Subjects with Autism Spectrum Disorders. *Mycopathologia* 182: 349002D363.
98. Finegold SM, Molitoris D, Song Y, Liu C, Vaisanen ML, et al. (2002) Gastrointestinal microflora studies in late-onset autism. *Clin Infect Dis* 35: S6-S16.
99. Onore C, Careaga M, Ashwood P (2012) The role of immune dysfunction in the pathophysiology of autism. *Brain Behav Immun* 26: 383-392.
100. Xu N, Li X, Zhong Y (2015) Inflammatory cytokines: Potential biomarkers of immunologic dysfunction in autism spectrum disorders. *Med Inflamm* 2015: 531518.
101. Siniscalco D, Bradstreet JJ, Antonucci N (2013) Therapeutic role of hematopoietic stem cells in autism spectrum disorder-related inflammation. *Front Immunol* 4: 140.
102. Ashwood P, Krakowiak P, Hertz-Pannier I, Hansen R, Pessah I, et al. (2011) Elevated plasma cytokines in autism spectrum disorders provide evidence of immune dysfunction and are associated with impaired behavioral outcome. *Brain Behav Immun* 25: 40-45.
- 103 Xie J, Huang L, Li X, Li H, Zhou Y, et al. (2017) Immunological cytokine profiling identifies TNF-α as a key molecule dysregulated in autistic children. *Oncotarget* 8: 82390-82398.
- 104 Jyonouchi H, Geng L, Davidow AL (2014) Cytokine profiles by peripheral blood monocytes are associated with changes in behavioral symptoms following immune insults in a subset of ASD subjects. *An*

- 124 Clark KB, Smith DC, Hassert DL, Browning RA, Naritoku DK, et al.  
(1998) Posttraining electrical stimulation of vagal afferents with  
concomitant vagal efferent inactivation enhances memory storage  
processes in the rat. *Neurobiol Learn Mem* 70: 364-373
- 125 Clark