

Notes on Neurodegenerative Criticism in Sleep

Adelino Canario*

Institute of Clinical Neurobiology, Innsbruck Medical University, Austria

Opinion

A recent study by S. N. H. N. (2022) [1] discusses the role of sleep in neurodegenerative diseases. The authors propose that sleep plays a critical role in the clearance of neurotoxic proteins, such as amyloid-beta and tau, from the brain. This process is mediated by the glymphatic system, which is activated during sleep. The study shows that sleep deprivation leads to an accumulation of these proteins, which in turn leads to neurodegeneration. This finding has important implications for the treatment of neurodegenerative diseases, as it suggests that improving sleep quality may be a potential therapeutic strategy. However, the study also highlights the need for further research to fully understand the mechanisms underlying the glymphatic system and its role in neurodegeneration.

328, A. PD, S. L. I. RAB1, L. T. 2. M. [4].

Multiple induration

Multiple induration (MS) is a condition characterized by the presence of multiple indurated areas on the skin. It is often associated with underlying systemic diseases, such as rheumatoid arthritis, systemic sclerosis, and Sjögren's syndrome. The pathogenesis of MS is not fully understood, but it is thought to be related to the deposition of immune complexes in the skin. Treatment of MS typically involves the use of corticosteroids and immunosuppressive agents. However, the response to treatment is often variable, and long-term management may be challenging. Further research is needed to better understand the underlying mechanisms of MS and to develop more effective treatment strategies.

C. (CJD) A. PRNP. C. (CJD) B.

Conclusion

S.

Parkinson's criticism

Parkinson's disease (PD) is a neurodegenerative disorder characterized by the loss of dopamine-producing neurons in the substantia nigra. This leads to motor symptoms such as tremor, rigidity, and bradykinesia. The pathogenesis of PD is complex and involves genetic and environmental factors. Current treatment strategies focus on dopamine replacement therapy, but there is a need for more effective and targeted treatments. Research in this area is ongoing, and it is hoped that a better understanding of the underlying mechanisms of PD will lead to improved outcomes for patients.

*Corresponding author: Adelino Canario, Institute of Clinical Neurobiology, Innsbruck Medical University, Austria, Phone: 1187421953230; E-mail: CanarioA@gmail.com
Received: 03-May-2022, Manuscript No. jcen-22-65168; **Editor assigned:** 05-May-2022, PreQC No. jcen-22-65168 (PQ); **Reviewed:** 12-May-2022, QC No. jcen-22-65168; **Revised:** 17-May-2022, Manuscript No. jcen-22-65168 (R); **Published:** 24-May-2022, DOI: 10.4172/jcen.1000150
Citation: Canario A (2022) Notes on Neurodegenerative Criticism in Sleep. J Clin Exp Neuroimmunol, 7: 150.
Copyright: © 2022 Canario A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

... I ... RAB1, ... L ... T ... 2 ... M ...

Acknowledgment

N ... , I ... M ... U ... , I ... C ...

Conflicts of Interest

... ..

References

1. Bodnar AG, et al (1998). Extension of lifespan by introduction of telomerase into normal human cells. *Science* 279(5349):349-352.
2. von Zglinicki T (2002). Oxidative stress shortens telomeres. *Trends Biochem Sci* 27:339-344.
3. Bartkova J, et al. (2006) Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature* 444(7119):633-637.
4. Eischen CM, Lozano G (2014). The Mdm network and its regulation of p53 activities: a rheostat of cancer risk. *Hum Mutat.* 35:728-737.
- 5.