

---

**\*Corresponding author:** Narenmandula, Department of Pharmacology, Toxicology and Biochemical Pharmaceutics, College of Pharmaceutical Sciences, Zhejiang University, China, E-mail: [narenman@zju.edu.cn](mailto:narenman@zju.edu.cn)

**Received** May 22, 2013; **Accepted** May 24, 2013; **Published** May 30, 2013

**Citation:** Narenmandula (2013) Novel Targets for Metal Compounds Inducing

comutagenicity. *Mutat Res* 478: 159-168.

8. Yá@Á ŠPÉÁ Ú^&\Á SÉÁ Š^Á VÓÁ ÇG€€GDÁ Ó@æ) \*^•Áž)Á \*^}^Á^Ç]!^••ž[}Á]! [,|^•Á [- @~ {æ}Á, à! [à|æ•c•š)Á!^•] [ ]•^k [áí~ {š!•^}šc^k!^æc {^}cÉÁÓæ!š) [ \*^}^•š• 23: 867-876.

9. Rehman K, Naranmandura H (2012) Arsenic metabolism and thioarsenicals. *Metallomics* 4: 881-892.

10. Naranmandura H, Xu S, Koike S, Pan LQ, Chen B, et al. (2012) The endoplasmic reticulum is a target organelle for trivalent dimethylarsinic acid (DMAIII)-induced cytotoxicity. *Toxicol Appl Pharmacol* 260: 241-249.