

Keywords:

Introduction

... () ... () ...
... 1-2 ... () ...
... () ...
... () ...
... 5 % ...
... 14% ...
... 3 ...
... 30 / ...
... 2 % ...
... 15 ...
... () ...
... 4 ...
... 5 ...
... 30% ... 50% ...
... 3 % ...
... 52% ...
... () ...

... 0 ... 0 ... 12 ... 0 ... () ... () - ...

... 20 ... 20 ... 100 ... 0 ... -12 ... 15 ...

... 20 ... 20 ... 100 ... 0 ... 0, ... 20, ... 110, (... 110, ...)).

... (...) ... 3 ...

... 5,000 ... 40 ... 2.4, ... 22. ... (...) ... 20, ... 2, ... 0 ...

... 0% ...

...

... 20-30% ...

... (50- 5% ...) ...

... (...) ... / ...

...

Conclusions

...

References

1. Tran DH, Maheshwari P, Nagaria Z, Patel HY, Verceles AC (2020) Ambulatory Status Is Associated With Successful Discharge Home in Survivors of Critical Illness. Respir Care 65: 1168-1173.
2. Piquet J, Brochard L, Isabey D, De Cremoux HT, Chang HK, et al. (1987) High frequency chest wall oscillation in patients with chronic air-fow obstruction. Am Rev Respir Dis 136: 1355-1359.
3. Gokdemir Y, Karadag Saygi E, Erdem E, Bayindir O, Ersu R, et al. (2014) Comparison of conventional pulmonary rehabilitation and high-frequency chest wall oscillation in primary ciliary dyskinesia. Pediatr Pulmonol 49: 611--2.01 d1orro0REVWU#0.9(ri61)7II. (1987)

9. González-Martín S, Becerro-de-Bengoa-Vallejo R, Angulo-Carrere MT, Iglesias ME, Martínez-Jiménez EM, et al. (2019) Effects of a visit prior to hospital admission on anxiety, depression and satisfaction of patients in an intensive care unit. *Intensive Criti Care Nurs* 54: 46-53.
10. Kho ME, Molloy AJ, Clarke FJ, Ajami D, McCaughan M, et al. (2016) TryCYCLE: a prospective study of the safety and feasibility of early in-bed cycling in mechanically ventilated patients. *PloS one* 11: 167561.
11. Rahimi RA, Skrzat J, Reddy DR, Zanni JM, Fan E, et al. (2013) Physical rehabilitation of patients in the intensive care unit requiring extracorporeal membrane oxygenation: a small case series. *Phys Ther* 93: 248-255.
12. Warwick WJ, Wielinski CL, Hansen LG (2004) Comparison of expectorated sputum after manual chest physical therapy and high-frequency chest compression.